

A Reliable Research Partner in Life Science and Medicine

Recombinant Human GM-CSF Protein(His Tag)

Catalog Number: PDMH100207

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Species Human

Source HEK293 Cells-derived Human GM-CSF proteins Ala18-Glu144, with an C-terminal

His

 Calculated MW
 13.9 kDa

 Observed MW
 20-30 kDa

 Accession
 P04141

Bio-activity Not validated for activity

Properties

Purity > 90% as determined by reducing SDS-PAGE.

Endotoxin < 1.0 EU/mg of the protein as determined by the LAL method

Storage Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -

80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of

reconstituted samples are stable at < -20°C for 3 months.

Shipping

This product is provided as lyophilized powder which is shipped with ice packs.

Formulation


Lyophilized from a 0.2 µm filtered solution in PBS with 5% Trehalose and 5%

Mannitol.

Reconstitution It is recommended that sterile water be added to the vial to prepare a stock solution

of 0.5 mg/mL. Concentration is measured by UV-Vis.

Data



SDS-PAGE analysis of Human GM-CSF proteins, 2 µg/lane of Recombinant Human GM-CSF proteins was resolved with SDS-PAGE under reducing conditions, showing bands at 13.9 KD

Background

For Research Use Only

 Toll-free: 1-888-852-8623
 Tel: 1-832-243-6086
 Fax: 1-832-243-6017

 Web: www.elabscience.com
 Email: techsupport@elabscience.com

Elabscience®

Elabscience Bionovation Inc.

A Reliable Research Partner in Life Science and Medicine

CSF2 (Colony Stimulating Factor 2) is a Protein Coding gene. Diseases associated with CSF2 include Mucositis and Candida Glabrata. Among its related pathways are RET signaling and IL-15 Signaling Pathways and their Primary Biological Effects in Different Immune Cell Types. GO annotations related to this gene include cytokine activity and granulocyte macrophage colony-stimulating factor receptor binding. The protein encoded by this gene is a cytokine that controls the production, differentiation, and function of granulocytes and macrophages. The active form of the protein is found extracellularly as a homodimer. This gene has been localized to a cluster of related genes at chromosome region 5q31, which is known to be associated with interstitial deletions in the 5q-syndrome and acute myelogenous leukemia. Other genes in the cluster include those encoding interleukins 4, 5, and 13.

For Research Use Only

 Toll-free: 1-888-852-8623
 Tel: 1-832-243-6086
 Fax: 1-832-243-6017

 Web: www.elabscience.com
 Email: techsupport@elabscience.com