
## **Elabscience**®

### **IDE Monoclonal Antibody**

#### catalog number: E-AB-22041

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description	
Reactivity	Human
Immunogen	Synthetic Peptide
Host	Mouse
Isotype	IgG
Clone	1H4
Purification	Protein A purification
Conjugation	Unconjugated
Buffer	Phosphate buffered solution, pH 7.4, containing 0.05% stabilizer, 0.5% protein protectant and 50% glycerol.
Applications	Recommended Dilution
WB	1:500-1:2000
IHC	1:50-300
IF	1:100-1:300
Data	



Negative Control

Western Blot analysis of 1) Hela, 2) HepG2 cells using IDE Monoclonal Antibody at dilution of 1:2000.

#### Observed-MW:118 kDa



Immunohistochemistry of paraffin-embedded Human liver cancer tissue using IDE Monoclonal Antibody at dilution of 1:200.

Immunofluorescence analysis of Human breast tissue using IDE Monoclonal Antibody at dilution of 1:200.

tore at -20°C Valid for 12 months. Avoid freeze / thaw cycles.
he product is shipped with ice pack, upon receipt, store it immediately at the emperature recommended.

Background

### For Research Use Only

Tel: 400-999-2100

# **Elabscience**®

This gene encodes a zinc metallopeptidase that degrades intracellular insulin, and thereby terminates insulins activity, as well as participating in intercellular peptide signalling by degrading diverse peptides such as glucagon, amylin, bradykinin, and kallidin. The preferential affinity of this enzyme for insulin results in insulin-mediated inhibition of the degradation of other peptides such as beta-amyloid. Deficiencies in this protein's function are associated with Alzheime r's disease and type 2 diabetes mellitus but mutations in this gene have not been shown to be causitive for these diseases. This protein localizes primarily to the cytoplasm but in some cell types localizes to the extracellular space, cell membrane, peroxisome, and mitochondrion. Alternative splicing results in multiple transcript variants encoding distinct isoforms. Additional transcript variants have been described but have not been experimentally verified.