## **Elabscience**®


## Recombinant Human AMBP/Alpha 1 Microglobulin Protein (His Tag)

## Catalog Number: PKSH033773

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Human HEK293 Cells-derived Human AMBP;Alpha 1 Microglobulin protein Gly20-Val203, with an C-terminal His 21.9 kDa 33 kDa P02760 Not validated for activity
with an C-terminal His 21.9 kDa 33 kDa P02760
21.9 kDa 33 kDa P02760
33 kDa P02760
P02760
Not validated for activity
> 95 % as determined by reducing SDS-PAGE.
< 1.0 EU per µg of the protein as determined by the LAL method.
Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80
°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of
reconstituted samples are stable at $< -20^{\circ}$ C for 3 months.
This product is provided as lyophilized powder which is shipped with ice packs.
Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4.
Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants
before lyophilization.
Please refer to the specific buffer information in the printed manual.
Please refer to the printed manual for detailed information.

Data



> 95 % as determined by reducing SDS-PAGE.

## Background

Protein AMBP belongs to the calycin superfamily and Lipocalin family. AMBP can be cleaved into three chains:  $\alpha$ -1microglobulin, inter- $\alpha$ -trypsin inhibitor light chain and trypstatin. AMBP is expressed by the liver and secreted in plasma.  $\alpha$ -1-microglobulin occurs in many physiological fluids including the plasma, urine, and cerebrospinal fluid. Inter- $\alpha$ trypsin inhibitor is present in the plasma and urine.  $\alpha$ -1-microglobulin occurs as a monomer and also in complexes with IgA and albumin, Inter- $\alpha$ -trypsin inhibitor inhibits trypsin, plasmin and lysosomal granulocytic elastase. Trypstatin act as a trypsin inhibitor, exists in a monomer forms and also occurs as a complex with tryptase in mast cells.