A Reliable Research Partner in Life Science and Medicine

Recombinant Human Fas/CD95/TNFRSF6 Protein (His Tag)

Catalog Number: PKSH033436

Note: Centrifuge before opening to ensure complete recovery of vial contents.

\mathbf{r}		crip					
	00	0	РΤ	n	ŤΤ		m
v	\mathbf{c}	v.		w	w	w	ш

Species Human

Source HEK293 Cells-derived Human Fas/CD95/TNFRSF6 protein Gln26-Asn173, with an C-

terminal His

Calculated MW 17.7 kDa
Observed MW 22-35 kDa
Accession P25445

Bio-activity Not validated for activity

Properties

Purity > 95 % as determined by reducing SDS-PAGE.

Endotoxin < 1.0 EU per µg of the protein as determined by the LAL method.

Storage Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80

°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of

reconstituted samples are stable at < -20°C for 3 months.

ShippingThis product is provided as lyophilized powder which is shipped with ice packs. **Formulation**Lyophilized from a 0.2 μm filtered solution of 20mM PB, 150mM NaCl, pH 7.4.


Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants

before lyophilization.

Please refer to the specific buffer information in the printed manual.

Reconstitution Please refer to the printed manual for detailed information.

Data



> 95 % as determined by reducing SDS-PAGE.

Background

FAS is a receptor and contains three TNFR-Cys repeats and one death domain. It has been shown that FAS is involved in the physiological regulation of programmed cell death, and has been implicated in the pathogenesis of various malignancies and diseases of the immune system. FADD (adapter molecule) recruits caspase-8 to the activated receptor, the resulting death-inducing signaling complex (DISC) performs caspase-8 proteolytic activation which initiates the subsequent cascade of caspases mediating apoptosis. FAS-mediated apoptosis may play a role in the induction of peripheral tolerance, in the antigen-stimulated suicide of mature T-cells, or both.

For Research Use Only