

Recombinant Human ILDR2 (C-6His)

Catalog Number: PKSH034056

Note: *Centrifuge before opening to ensure complete recovery of vial contents.*


Description

Species	Human
Source	HEK293 Cells-derived Human ILDR2 protein Leu21-Glu186, with an C-terminal His
Calculated MW	19.5 kDa
Observed MW	45-55 kDa
Accession	Q71H61
Bio-activity	Not validated for activity

Properties

Purity	> 95 % as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU per µg of the protein as determined by the LAL method.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4. Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization.
Reconstitution	Please refer to the specific buffer information in the printed manual. Please refer to the printed manual for detailed information.

Data



> 95 % as determined by reducing SDS-PAGE.

Background

ILDR2 is a member of the B7-like family of proteins that regulate T cell activity, is also a known endoplasmic reticulum molecule that regulates lipid homeostasis. The human ILDR2 luminal domain shares a 99% and 98% homology with the mouse and rat respectively. The human gene encoding ILDR2 is located in a region on Chr1q23–25 that has been associated with type 2 diabetes. ILDR2 plays critical roles in hepatic clearance of lipoproteins and in lipid homeostasis. ILDR2 regulates human dendritic cells (DC2 cells, a subpopulation of polarized DCs that promotes Th2 differentiation). Recent publications reported that ILDR2 displayed negative regulatory functions on human and mouse T cells in various experimental systems. Fusion protein of ILDR2 luminal domain with an Fc fragment, displays therapeutic effects in collagen-induced arthritis (CIA), a mouse model of rheumatoid arthritis (RA). ILDR2 represents a novel B7-like ligand that exerts negative immune modulation via interaction with a putative counterpart receptor expressed on activated T cells.

For Research Use Only

Toll-free: 1-888-852-8623
Web: www.elabscience.com

Tel: 1-832-243-6086
Email: techsupport@elabscience.com

Fax: 1-832-243-6017