Recombinant Mouse MEK1/MAP2K1/MKK1 Protein

Catalog Number: PKSM040746

Note: *Centrifuge before opening to ensure complete recovery of vial contents.*


Description

Species	Mouse
Source	Baculovirus-Insect Cells-derived Mouse MEK1/MAP2K1/MKK1 protein Met1-Ile393
Calculated MW	43.6 kDa
Observed MW	45 kDa
Accession	NP_032953.1
Bio-activity	Kinase activity untested

Properties

Purity	> 90% as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU per μg of the protein as determined by the LAL method.
Storage	Store at < -20°C, stable for 6 months. Please minimize freeze-thaw cycles.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Supplied as sterile 20 mM Tris, 500 mM NaCl, 10% glycerol, pH 8.0 Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.

Data



> 90 % as determined by reducing SDS-PAGE.

Background

MEK1, also known as MAP2K1 and MKK1, is a member of the dual specificity protein kinase family, which acts as a mitogen-activated protein (MAP) kinase kinase. MAP kinases, also known as extracellular signal-regulated kinases (ERKs), act as an integration point for multiple biochemical signals. MEK1 is widely expressed, with extremely low levels in brain. It lies upstream of MAP kinases and stimulates the enzymatic activity of MAP kinases upon wide variety of extra- and intracellular signals. As an essential component of MAP kinase signal transduction pathway, MEK1 is involved in many cellular processes such as proliferation, differentiation, transcription regulation and development. Binding extracellular ligands such as growth factors, cytokines and hormones to their cell-surface receptors activates RAS and this initiates RAF1 activation. RAF1 then further activates the dual-specificity protein kinases MAP2K1 and MEK2. MEK1 has been shown to export PPARG from the nucleus. The MAPK cascade is also involved in the regulation of endosomal dynamics, including lysosome processing and endosome cycling through the perinuclear recycling compartment (PNRC), as well as in the fragmentation of the Golgi apparatus during mitosis. MKK1 catalyzes the concomitant phosphorylation of a threonine and a tyrosine residue in a Thr-Glu-Tyr sequence located in MAP kinases. Defects in MEK1 can cause cardiofaciocutaneous syndrome.

Toll-free: 1-888-852-8623
Tel: 1-832-243-6086
Fax: 1-832-243-6017
Web: www.elabscience.com
Email: techsupport@elabscience.com