

A Reliable Research Partner in Life Science and Medicine

# Recombinant Human SIRPalpha/CD172a Protein (His Tag)

Catalog Number: PDEH101043

Note: Centrifuge before opening to ensure complete recovery of vial contents.

#### Description

Species Human

Source E.coli-derived Human SIRPA protein Glu31-Ser350, with an N-terminal His & C-

terminal His

 Calculated MW
 35.1 kDa

 Observed MW
 39 kDa

 Accession
 P78324-2

Bio-activity Not validated for activity

### **Properties**

**Purity** > 95% as determined by reducing SDS-PAGE.

Endotoxin < 10 EU/mg of the protein as determined by the LAL method

Storage Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -

80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of

reconstituted samples are stable at < -20°C for 3 months.

Shipping

This product is provided as lyophilized powder which is shipped with ice packs.

Formulation


Lyophilized from a 0.2 µm filtered solution in PBS with 5% Trehalose and 5%

Mannitol.

Reconstitution It is recommended that sterile water be added to the vial to prepare a stock solution

of 0.5 mg/mL. Concentration is measured by UV-Vis.

## Data



SDS-PAGE analysis of Human SIRPalpha/CD172a proteins, 2 µg/lane of Recombinant Human SIRPalpha/CD172a proteins was resolved with SDS-PAGE under reducing conditions, showing bands at 39 kDa.

## **Background**

For Research Use Only

 Toll-free: 1-888-852-8623
 Tel: 1-832-243-6086
 Fax: 1-832-243-6017

 Web: www.elabscience.com
 Email: techsupport@elabscience.com

Rev. V1.4

# Elabscience®

#### Elabscience Bionovation Inc.

A Reliable Research Partner in Life Science and Medicine

Signal Regulatory Protein  $\alpha$  (SIRP $\alpha$ ) is a monomeric approximately 90 kD type I transmembrane glycoprotein. The 504 amino acid human SIRP $\alpha$  contains two Ig-like C1-type domains and one Ig-like V-type domain. SIRP $\alpha$  can express in various tissues, mainly on brain and myeloid cells, including macrophages, neutrophils, dendritic and Langerhans cells. It also can detect in neurons, smooth muscle and endothelial cells. SIRPA is an immunoglobulin-like cell surface receptor for CD47. SIRP $\alpha$  acts as docking protein and induces translocation of PTPN6, PTPN11 and other binding partners from the cytosol to the plasma membrane. SIRP $\alpha$  shows adhesion of cerebellar neurons, neurite outgrowth and glial cell attachment. SIRP $\alpha$  engagement generally produces a negative regulatory signal, it may mediate negative regulation of phagocytosis, mast cell activation and dendritic cell activation

For Research Use Only

 Toll-free: 1-888-852-8623
 Tel: 1-832-243-6086
 Fax: 1-832-243-6017

 Web: www.elabscience.com
 Email: techsupport@elabscience.com