







BMP5 Polyclonal Antibody

catalog number: E-AB-62013

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description	
Reactivity	Human;Mouse;Rat
Immunogen	Recombinant fusion protein of human BMP5 (NP_066551.1).
Host	Rabbit
Isotype	Rabbit IgG
Purification	Affinity purification
Buffer	Phosphate buffered solution, pH 7.4, containing 0.05% stabilizer and 50% glycerol.
Applications	Recommended Dilution
WB	1:500-1:2000
IHC	1:50-1:100

Data	
 Western blot analysis of extracts of various cell lines using BMP5 Polyclonal Antibody at dilution of 1:1000. Observed-MW:52 kDa Calculated-MW:47 kDa/51 kDa	 Immunohistochemistry of paraffin-embedded Rat lung using BMP5 Polyclonal Antibody at dilution of 1:100 (40x lens).
 Immunohistochemistry of paraffin-embedded Rat brain using BMP5 Polyclonal Antibody at dilution of 1:100 (40x lens).	 Immunohistochemistry of paraffin-embedded Human prostate using BMP5 Polyclonal Antibody at dilution of 1:100 (40x lens).
 Immunohistochemistry of paraffin-embedded Human esophagus using BMP5 Polyclonal Antibody at dilution of 1:100 (40x lens).	 Immunohistochemistry of paraffin-embedded Mouse lung using BMP5 Polyclonal Antibody at dilution of 1:100 (40x lens).

Preparation & Storage

For Research Use Only

Toll-free: 1-888-852-8623
Web: www.elabscience.com

Tel: 1-832-243-6086
Email: techsupport@elabscience.com

Fax: 1-832-243-6017

Rev. V1.8

Storage

Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.

Shipping

The product is shipped with ice pack, upon receipt, store it immediately at the temperature recommended.

Background

This gene encodes a member of the bone morphogenetic protein family which is part of the transforming growth factor-beta superfamily. The superfamily includes large families of growth and differentiation factors. Bone morphogenetic proteins were originally identified by an ability of demineralized bone extract to induce endochondral osteogenesis in vivo in an extraskeletal site. These proteins are synthesized as prepropeptides, cleaved, and then processed into dimeric proteins. This protein may act as an important signaling molecule within the trabecular meshwork and optic nerve head, and may play a potential role in glaucoma pathogenesis. This gene is differentially regulated during the formation of various tumors.

For Research Use Only

Toll-free: 1-888-852-8623

Web: www.elabscience.com

Tel: 1-832-243-6086

Email: techsupport@elabscience.com

Fax: 1-832-243-6017

Rev. V1.8