Elabscience®

Human DPP4 Antibody Pair Set

Catalog No.E-KAB-0022ApplicationsELISASynonymsCD26, DPPIV, ADABP, ADCP2, TP103, Adenosine Deaminase Complexing Protein 2

Kit components & Storage

Title	Specifications	Storage
Human DPP4 Capture Antibody	1 vial, 100 µ g	Store at -20° C for one year.
		Avoid freeze / thaw cycles.
Human DPP4 Detection Antibody	1 vial, 50 μL	Store at -20°C for one year.
(Biotin)		Avoid freeze / thaw cycles.

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Product Information

Items		Characteristic (E-KAB-0022)	
		Human DPP4 Capture Antibody	Human DPP4 Detection Antibody
			(Biotin)
Immunogen	Immunogen	Recombinant Human DPP4 protein	Recombinant Human DPP4 protein
Information	Swissprot	P27487	
Product details	Reactivity	Human	Human
	Host	Rat	Goat
	Conjugation	Unconjugated	Biotin
	Concentration	0.5mg/mL	/
	Buffer	PBS with 0.04% Proclin 300, 50%	PBS with 0.04% Proclin 300, 1%
		glycerol, pH 7.4	protective protein, 50% glycerol, pH
			7.4
	Purify	Protein A or G	Antigen Affinity
	Specificity	Detects Human DPP4 in ELISAs.	

For Research Use Only

Elabscience®

Applications

Human DPP4 Sandwich ELISA Assay:

Recommended	Reagent	Images
Concentration/Dilution		
0.5-4µg/mL	Human DPP4 Capture Antibody	
		Aig 1
1:1000-1:10000	Human DPP4 Detection Antibody	Optical Density
	(Biotin)	O 0.1
		0.01
-	Concentration/Dilution 0.5-4µg/mL	Concentration/Dilution 0.5-4µg/mL Human DPP4 Capture Antibody 1:1000-1:10000 Human DPP4 Detection Antibody

Note: This standard curve is only for demonstration purposes. A standard curve should be generated for each assay!

Background

The protein encoded by this gene is identical to adenosine deaminase complexing protein-2, and to the T-cell activation antigen CD26. It is an intrinsic membrane glycoprotein and a serine exopeptidase that cleaves X-proline dipeptides from the N-terminus of polypeptides.

DPP4 (Dipeptidyl Peptidase 4) is a Protein Coding gene. Diseases associated with DPP4 include Nasopharyngitis and Mental Depression. Among its related pathways are Collagen chain trimerization and Incretin synthesis, secretion, and inactivation. GO annotations related to this gene include protein homodimerization activity and receptor binding. An important paralog of this gene is FAP.

For Research Use Only