Elabscience®

ELISA

Mouse Cys-C Antibody Pair Set

| Catalog No. | E-KAB-0073 | Applications |
|-------------|---|--------------|
| Synonyms | CST3, Cystatin 3, ARMD11, Post-Gamma-Globulin | |

Kit components & Storage

| Title | Specifications | Storage |
|--------------------------------|-----------------|--|
| Mouse Cys-C Capture Antibody | 1 vial, 100 µ g | Store at -20° C for one year. |
| | | Avoid freeze / thaw cycles. |
| Mouse Cys-C Detection Antibody | 1 vial, 50 μL | Store at -20° C for one year. |
| (Biotin) | | Avoid freeze / thaw cycles. |

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Product Information

| Items | | Characteristic (E-KAB-0073) | |
|-----------------|---------------|--|---|
| | | Mouse Cys-C Capture Antibody | Mouse Cys-C Detection Antibody (Biotin) |
| Immunogen | Immunogen | Recombinant Mouse Cys-C protein | Recombinant Mouse Cys-C protein |
| Information | Swissprot | P21460 | |
| Product details | Reactivity | Mouse | Mouse |
| | Host | Goat | Goat |
| | Conjugation | Unconjugated | Biotin |
| | Concentration | 0.5mg/mL | / |
| | Buffer | PBS with 0.04% Proclin 300, 50% glycerol, pH 7.4 | PBS with 0.04% Proclin 300, 1% protective protein, 50% glycerol, pH 7.4 |
| | Purify | Antigen Affinity | Antigen Affinity |
| | Specificity | Detects Mouse Cys-C in ELISAs. | |

For Research Use Only

Elabscience®

Applications

Mouse Cys-C Sandwich ELISA Assay:

| | Recommended | Reagent | Images |
|--------------------|------------------------|--|---|
| | Concentration/Dilution | | |
| ELISA | 0.5-4µg/mL | Mouse Cys-C Capture Antibody | |
| Capture | | | |
| ELISA Detection | 1:1000-1:10000 | Mouse Cys-C Detection Antibody (Biotin) | Optical Density |
| | | | 0.01 100 1000 10000 10000 Mouse Cys-C concentration(pg/mL) |

Note: This standard curve is only for demonstration purposes. A standard curve should be generated for each assay!

Background

Cystatin C is a member of family 2 of the Cystatin superfamily. It is involved in processes such as tumor invasion and metastasis, inflammation and some neurological diseases. It inhibits many cysteine proteases such as papain and cathepsins B,H,K,L,and S.

For Research Use Only