Recombinant Mouse TNFR1/TNFRSF1A Protein (His Tag)

Catalog Number: PKSM040648

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description	
Species	Mouse
Source	HEK293 Cells-derived Mouse TNFR1/TNFRSF1A protein Met 1-Ala 212, with an C-
	terminal His
Calculated MW	21.8 kDa
Accession	NP_035739.2
Bio-activity	1. Immobilized human TNFa at 10 μ g/mL (100 μ l/well) can bind biotinylated mouse
	TNFRSF1A-his. The EC ₅₀ of biotinylated mouse TNFRSF1A-his is 0.28 μ g/mL. 2. Immobilized mouse TNFa at 10 μ g/mL (100 μ l/well) can bind biotinylated mouse
	TNFRSF1A-his. The EC ₅₀ of biotinylated mouse TNFRSF1A-his is 0.13 μ g/mL. 3. Measured by its ability to inhibit the TNF α -mediated cytotoxicity in the L929 mouse
	fibrosarcoma cells in the presence of the metabolic inhibitor actinomycin D. The ED_{50} for this effect is typically 0.5-1. 5 µg/mL in the presence of 0.1 ng/mL recombinant
	mouse TNFa.
Properties	
Purity	> 96 % as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU per µg of the protein as determined by the LAL method.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80
	°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of
	reconstituted samples are stable at $<$ -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from sterile PBS, pH 7.4
	Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants
	before lyophilization.
	Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.

Data



> 96 % as determined by reducing SDS-PAGE.

Background

For Research Use Only

Elabscience®

The cluster of differentiation (CD) system is commonly used as cell markers in immunophynotyping. Different kinds of cells in the immune system can be identified through the surface CD molecules which associating with the immune function of the cell. There are more than 320 CD unique clusters and subclusters have been identified. Some of the CD molecules serve as receptors or ligands important to the cell through initiating a signal cascade which then alter the behavior of the cell. Some CD proteins do not take part in cell signal process but have other functions such as cell adhesion. CD120a (cluste of differentiation 120a), also known as TNFR1 / TNFRSF1A, is a member of CD family, tumor necrosis factor receptor superfamily. CD120a is one of the most primary receptors for the tumor necrosis factor-alpha. It has been shown to be localized to both plasma membrane lipid rafts and the trans golgi complex with the help of the death domain (DD). CD120a can activate the transcription factor NF-κB, mediate apoptosis, and regulate inflammation processes.