

A Reliable Research Partner in Life Science and Medicine

SLC27A2 Polyclonal Antibody

catalog number: E-AB-60173

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Reactivity Human; Mouse

Immunogen Recombinant fusion protein of human SLC27A2 (NP 001153101.1).

Host Rabbit
Isotype IgG


Purification Affinity purification

Buffer Phosphate buffered solution, pH 7.4, containing 0.05% stabilizer and 50% glycerol.

Applications Recommended Dilution


WB 1:500-1:2000 **IHC** 1:50-1:200

Data




Western blot analysis of extracts of HepG2 cells using SLC27A2 Polyclonal Antibody.

Observed-MW:70 kDa Calculated-MW:64 kDa/70 kDa



Immunohistochemistry of paraffin-embedded Human gastric cancer using SLC27A2 Polyclonal Antibody at dilution of 1:100 (40x lens).



Immunohistochemistry of paraffin-embedded Human liver using SLC27A2 Polyclonal Antibody at dilution of 1:100 (40x lens).

Preparation & Storage

Storage Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.

Shipping The product is shipped with ice pack,upon receipt, store it immediately at the

temperature recommended.

Background

For Research Use Only

Elabscience Bionovation Inc.

A Reliable Research Partner in Life Science and Medicine

Elabscience®

The protein encoded by this gene is an isozyme of long-chain fatty-acid-coenzyme A ligase family. Although differing in substrate specificity, subcellular localization, and tissue distribution, all isozymes of this family convert free long-chain fatty acids into fatty acyl-CoA esters, and thereby play a key role in lipid biosynthesis and fatty acid degradation. This isozyme activates long-chain, branched-chain and very-long-chain fatty acids containing 22 or more carbons to their CoA derivatives. It is expressed primarily in liver and kidney, and is present in both endoplasmic reticulum and peroxisomes, but not in mitochondria. Its decreased peroxisomal enzyme activity is in part responsible for the biochemical pathology in X-linked adrenoleukodystrophy. Alternatively spliced transcript variants encoding different isoforms have been found for this gene.

Fax: 1-832-243-6017