

Recombinant Mouse Fetuin B Protein(Fc Tag)

Catalog Number: PDMM100140

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Species	Mouse
Source	Mammalian-derived Mouse Fetuin B proteins Arg19-Pro388,with an C-terminal Fc
Mol_Mass	65.6 kDa
Accession	Q9QXC1
Bio-activity	Not validated for activity

Properties

Purity	> 90% as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU/mg of the protein as determined by the LAL method
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80 °C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from a 0.2 µm filtered solution in PBS with 5% Trehalose and 5% Mannitol.
Reconstitution	It is recommended that sterile water be added to the vial to prepare a stock solution of 0.5 mg/mL. Concentration is measured by UV-Vis.

Data



SDS-PAGE analysis of Mouse Fetuin B proteins , 2µg/lane of

Recombinant Mouse Fetuin B proteins was resolved with SDS-PAGE under reducing conditions , showing bands at 80

KD

Background

Fetuin-B, also known as Fetuin-like protein IRL685 and FETUB, is a secreted protein which belongs to the fetuin family. Fetuin-B / FETUB contains two cystatin domains. Fetuin-B is a member of the fetuin family, part of the cystatin superfamily of cysteine protease inhibitors. Fetuins have been implicated in several diverse functions, including osteogenesis and bone resorption. Fetuin-A has been identified as a major protein during fetal life and is also involved in important functions such as protease inhibitory activities and development-associated regulation of calcium metabolism and osteogenesis. Fetuin-A is a key partner in the recovery phase of an acute inflammatory response. Fetuin-B / FETUB is found at least in Human and rodents. It is unambiguously a paralogue of Fetuin-A. Fetuin-A and Fetuin-B exhibit significant differences at the amino acid sequence level, notably including variations with respect to the archetypal fetuin-specific signature.

For Research Use Only