

APC Anti-Human IL-10 Antibody[JES3-9D7]

Catalog Number: GFH1198E

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description


Reactivity	Human
Host	Rat
Isotype	Rat IgG1, κ
Clone No.	JES3-9D7
Isotype Control	APC Rat IgG1, κ Isotype Control[HRPN] [Product E-AB-F09822E]
Conjugation	APC
Conjugation Information	APC is designed to be excited by the Red (627-640 nm) laser and detected using an optical filter centered near 660 nm (e.g., a 660/20 nm bandpass filter).
Storage Buffer	Phosphate buffered solution, pH 7.2, containing 0.09% sodium azide.

Applications

Recommended usage

FCM	Each lot of this antibody is quality control tested by flow cytometric analysis. The amount of the reagent is suggested to be used 5 μL of antibody per test (million cells in 100 μL staining volume or per 100 μL of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.
------------	--

Data



HEK293T cells transiently transfected with pcDNA3.1 plasmid encoding Human IL-10 gene are stained with APC Anti-Human IL-10 Antibody (filled gray histogram) or APC Rat IgG1, κ Isotype Control (empty black histogram).

Preparation & Storage

Storage	Keep as concentrated solution. This product can be stored at 2-8°C for 12 months. Please protected from prolonged exposure to light and do not freeze.
Shipping	Ice bag

Antigen Information

Alternate Names	CSIF;Cytokine synthesis inhibitory factor;IL-10;Interleukin-10
Uniprot ID	P22301
Gene ID	3586

For Research Use Only

Background

IL-10 was originally described as Cytokine Synthesis Inhibitory Factor (CSIF) by virtue of its ability to inhibit cytokine production by Th1 clones. IL-10 shares over 80% sequence homology with the Epstein-Barr virus protein BCRF1. The biological activities of IL-10 include inhibition of macrophage-mediated cytokine synthesis, suppression of the delayed type hypersensitivity response, and stimulation of the Th2 cell response, which results in elevated antibody production.