



A Reliable Research Partner in Life Science and Medicine

Purified Anti-Human CD19 Antibody[4G7]

catalog number: E-AB-F11270P

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Reactivity Human

Immunogen Recombinant Human CD19 protein

Host Mouse

Isotype Mouse IgG1, κ

Clone 4G7

Purification >98%, Protein A/G purified


Buffer Phosphate-buffered solution, pH 7.2, containing 0.05% non-protein stabilizer.

Dialyze to completely remove the stabilizer prior to labeling.

Applications Recommended Dilution

FCM $2 \mu g/mL(0.5 \times 10^6 - 1 \times 10^6 \text{ cells})$

Data



Human peripheral blood lymphocytes were stained with 0.2 μg Purified Anti-Human CD19 Antibody[4G7] (Right) and 0.2 μg Mouse IgG1, κ Isotype Control (Left), followed by APC-conjugated Goat Anti-Mouse IgG Secondary Antibody, then anti-Human CD22 PE-conjugated Monoclonal Antibody.

Preparation & Storage

Storage Store at 4°C valid for 12 months or -20°C valid for long term storage, avoid freeze /

thaw cycles.

Shipping Ice bag

Background

CD19 protein is a 95 kDa transmembrane glycoprotein that plays a central role in B cell activation and humoral immune responses . CD19 is expressed throughout B cell development from pre-B cells through mature B cells, and it is commonly used as a B cell lineage marker . It is required for the responsiveness of mature B cell to antigen stimulation, germinal center development, and antibody affinity maturation . The CD19 protein associates with the B cell antigen receptor (BCR), CD81, CD38, CD21, CD22, and IFITM1/CD225/Leu13. These associations enable CD19 to amplify B cell signaling and reduce the threshold for antigen stimulation through the BCR.

For Research Use Only

 Toll-free: 1-888-852-8623
 Tel: 1-832-243-6086
 Fax: 1-832-243-6017

 Web: www.elabscience.com
 Email: techsupport@elabscience.com
 Rev. V1.6