A Reliable Research Partner in Life Science and Medicine

# Recombinant Human CD96 Protein (His Tag)

Catalog Number: PDMH100422

Note: Centrifuge before opening to ensure complete recovery of vial contents.

#### Description

Species Human

Source HEK293 Cells-derived Human CD96 protein Met1-Met519, with an C-terminal His

 Calculated MW
 57.0 kDa

 Observed MW
 58 kDa

 Accession
 P40200

Bio-activity Not validated for activity

#### **Properties**

**Purity** > 95% as determined by reducing SDS-PAGE.

Endotoxin < 1.0 EU/mg of the protein as determined by the LAL method

Storage Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -

80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of

reconstituted samples are stable at < -20°C for 3 months.


ShippingThis product is provided as lyophilized powder which is shipped with ice packs.FormulationLyophilized from a 0.2 μm filtered solution in PBS with 5% Trehalose and 5%

Mannitol

**Reconstitution** It is recommended that sterile water be added to the vial to prepare a stock solution

of 0.5 mg/mL. Concentration is measured by UV-Vis.

#### Data



SDS-PAGE analysis of Human CD96 proteins, 2 µg/lane of Recombinant Human CD96 proteins was resolved with SDS-PAGE under reducing conditions, showing bands at 58 kDa.

### **Background**

The cluster of differentiation (CD) system is commonly used as cell markers in immunophynotyping. Different kinds of cells in the immune system can be identified through the surface CD molecules which associating with the immune function of the cell. The CD155 ligand CD96 is a member of the lg superfamily. It&#39,s a immunoglobulin-like protein tentatively allocated to the repertoire of human NK receptors. NK cells recognize poliovirus receptor (PVR), anectins and nectin-like protein family member serve to mediate cell-cell adhesion, cell migration, with the presence of an additional receptor, CD96. CD96 promotes NK cell adhesion to target cells expressing PVR, stimulates cytotoxicity of activated NK cells, and mediates acquisition of PVR from target cells.

## For Research Use Only

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017

Web: www.elabscience.com Email: techsupport@elabscience.com