Elabscience®

Human MMP-3 Antibody Pair Set

Catalog No.E-KAB-0054ApplicationsSynonymsMMP3, CHDS6, SL-1, STMY, STMY1, STR1, Stromelysin-1

ELISA

Kit components & Storage

Title	Specifications	Storage
Human MMP-3 Capture Antibody	1 vial, 100 µ g	Store at -20° C for one year.
		Avoid freeze / thaw cycles.
Human MMP-3 Detection Antibody	1 vial, 50 μL	Store at -20°C for one year.
(Biotin)		Avoid freeze / thaw cycles.

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Product Information

Items		Characteristic (E-KAB-0054)	
		Human MMP-3 Capture Antibody	Human MMP-3 Detection Antibody
			(Biotin)
Immunogen	Immunogen	Recombinant Human MMP-3 protein	Recombinant Human MMP-3 protein
Information	Swissprot	P08254	
Product details	Reactivity	Human	Human
	Host	Goat	Goat
	Conjugation	Unconjugated	Biotin
	Concentration	0.5mg/mL	/
	Buffer	PBS with 0.04% Proclin 300, 50%	PBS with 0.04% Proclin 300, 1%
		glycerol, pH 7.4	protective protein, 50% glycerol, pH
			7.4
	Purify	Antigen Affinity	Antigen Affinity
	Specificity	Detects Human MMP-3 in ELISAs.	

For Research Use Only

Elabscience®

Applications

Human MMP-3 Sandwich ELISA Assay:

	Recommended	Reagent	Images
	Concentration/Dilution		
ELISA	0.5-4µg/mL	Human MMP-3 Capture Antibody	
Capture			
ELISA Detection	1:1000-1:10000	Human MMP-3 Detection Antibody (Biotin)	Optical Density
			0.01 0.1 1 10 100 0.01 0.1 1 10 100 Human MMP-3 concentration(ng/mL)

Note: This standard curve is only for demonstration purposes. A standard curve should be generated for each assay!

Background

Proteins of the matrix metalloproteinase (MMP) family are involved in the breakdown of extracellular matrix in normal physiological processes, such as embryonic development, reproduction, and tissue remodeling, as well as in disease processes, such as arthritis and metastasis. Most MMP's are secreted as inactive proproteins which are activated when cleaved by extracellular proteinases. This gene encodes an enzyme which degrades fibronectin, laminin, collagens III, IV, IX, and X, and cartilage proteoglycans. The enzyme is thought to be involved in wound repair, progression of atherosclerosis, and tumor initiation. The gene is part of a cluster of MMP genes which localize to chromosome 11q22.3.