


## YWHAG Polyclonal Antibody

catalog number: E-AB-12587

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.


| Description  |  |
|--------------|--|
| Reactivity   | Human;Mouse;Rat  |
| Immunogen    | Synthetic peptide of human YWHAG   |
| Host         | Rabbit   |
| Isotype      | IgG  |
| Purification | Affinity purification  |
| Conjugation  | Unconjugated   |
| Buffer       | Phosphate buffered solution, pH 7.4, containing 0.05% stabilizer and 50% glycerol. |
| Applications | Recommended Dilution   |
| WB           | 1:1000-1:5000  |
| IHC          | 1:50-1:200   |

## Data




Western Blot analysis of 293T and Jurkat cell using YWHAG Polyclonal Antibody at dilution of 1:800

Calculated-MW:28 kDa



Immunohistochemistry of paraffin-embedded Human brain using YWHAG Polyclonal Antibody at dilution of 1:50



Immunohistochemistry of paraffin-embedded Human lung cancer using YWHAG Polyclonal Antibody at dilution of 1:50

| Preparation & Storage |  |
|-----------------------|--|
| Storage               | Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.  |
| Shipping              | The product is shipped with ice pack, upon receipt, store it immediately at the temperature recommended. |

**Background**

This gene product belongs to the 14-3-3 family of proteins which mediate signal transduction by binding to phosphoserine-containing proteins. This highly conserved protein family is found in both plants and mammals, and this protein is 100% identical to the rat ortholog. It is induced by growth factors in human vascular smooth muscle cells, and is also highly expressed in skeletal and heart muscles, suggesting an important role for this protein in muscle tissue. It has been shown to interact with RAF1 and protein kinase C, proteins involved in various signal transduction pathways.

## For Research Use Only