Recombinant Mouse GM-CSF protein (His Tag)

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Catalog Number: PDEM100256



Description Species Mouse Source E.coli-derived Mouse GM-CSF protein Ala18-Lys141, with an N-terminal His Mol Mass 13.5 kDa P01587 Accession **Bio-activity** Not validated for activity **Properties** Purity >95% as determined by reducing SDS-PAGE. Endotoxin < 10 EU/mg of the protein as determined by the LAL method Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80 Storage °C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at $< -20^{\circ}$ C for 3 months. Shipping This product is provided as lyophilized powder which is shipped with ice packs. Lyophilized from a 0.2 µm filtered solution in PBS with 5% Trehalose and 5% Formulation Mannitol. Reconstitution It is recommended that sterile water be added to the vial to prepare a stock solution of 0.5 mg/mL. Concentration is measured by UV-Vis. Data

KDa	М	R
80		
60		
40		
30		
20		
12	_	

> 95 % as determined by reducing SDS-PAGE.

Background

Granulocyte-Macrophage Colony Stimulating Factor (GM-CSF) was initially characterized as a growth factorthat can support the in vitro colony formation of granulocyte-macrophage progenitors. It is produced by anumber of different cell types (including activated T cells, B cells, macrophages, mast cells, endothelial cellsand fibroblasts) in response to cytokine of immune and inflammatory stimuli. Besides granulocyte-macrophageprogenitors, GM-CSF is also a growth factor for erythroid, megakaryocyte and eosinophil progenitors. Onmature hematopoietic, monocytes/ macrophages and eosinophils. GM-CSF has a functional role on nonhematopoitic cells. It can induce human endothelial cells to migrate and proliferate. Additionally, GM-CSF canalso stimulate the proliferation of a number of tumor cell lines, including osteogenic sarcoma, carcinoma andadenocarcinoma cell lines.

For Research Use Only