

Recombinant Human/Mouse USP46 Protein (SUMO Tag)

Catalog Number: PKSH030765

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Species	Human
Source	Baculovirus-Insect Cells-derived Human USP46 protein Met 1-Glu366, with an N-terminal SUMO
Calculated MW	55.2 kDa
Observed MW	64 kDa
Accession	NP_808229.1
Bio-activity	Not validated for activity

Properties

Purity	> 90 % as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU per µg of the protein as determined by the LAL method.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from sterile 20mM Tris, 500mM NaCl, pH 7.4, 10% glycerol Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.

Data



> 90 % as determined by reducing SDS-PAGE.

Background

USP46 belongs to the peptidase C19 family; USP12/USP46 subfamily. Deubiquitinating enzymes (DUBs) are a large group of proteases which are also commonly referred to as deubiquitinating peptidases; deubiquitinating isopeptidases; deubiquitinases; ubiquitin proteases; ubiquitin hydrolyases; ubiquitin isopeptidases; or Dubs. They regulate ubiquitin-dependent metabolic pathways by cleaving ubiquitin-protein bonds. They also may act as negative and positive regulators of the ubiquitin system. Besides ubiquitin recycling; they are also involved in processing of ubiquitin precursors; in proofreading of protein ubiquitination and in disassembly of inhibitory ubiquitin chains. USP46 is a deubiquitinating enzyme that plays a role in behavior; possibly by regulating GABA action. It may act by mediating the deubiquitination of GAD1/GAD67. USP46 has almost no deubiquitinating activity by itself and requires the interaction with WDR48 to have a high activity and it is not involved in deubiquitination of monoubiquitinated FANCD2.

For Research Use Only

Toll-free: 1-888-852-8623
Web: www.elabscience.com

Tel: 1-832-243-6086
Email: techsupport@elabscience.com

Fax: 1-832-243-6017