

# Recombinant Mouse CD27/TNFRSF7 Protein(Fc Tag)

Catalog Number: PDMM100150



**Note:** Centrifuge before opening to ensure complete recovery of vial contents.


## Description

<b>Species</b>	Mouse
<b>Source</b>	Mammalian-derived Mouse CD27/TNFRSF7 proteins Pro24-Arg182,with an C-terminal Fc
<b>Mol_Mass</b>	42.4 kDa
<b>Accession</b>	P41272
<b>Bio-activity</b>	Not validated for activity

## Properties

<b>Purity</b>	> 90% as determined by reducing SDS-PAGE.
<b>Endotoxin</b>	< 1.0 EU/mg of the protein as determined by the LAL method
<b>Storage</b>	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80 °C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
<b>Shipping</b>	This product is provided as lyophilized powder which is shipped with ice packs.
<b>Formulation</b>	Lyophilized from a 0.2 µm filtered solution in PBS with 5% Trehalose and 5% Mannitol.
<b>Reconstitution</b>	It is recommended that sterile water be added to the vial to prepare a stock solution of 0.5 mg/mL. Concentration is measured by UV-Vis.

## Data



SDS-PAGE analysis of Mouse CD27/TNFRSF7 proteins ,  
2µg/lane of Recombinant Mouse CD27/TNFRSF7 proteins  
was resolved with SDS-PAGE under reducing conditions ,  
showing bands at 60 KD

## Background

CD27 , also known as TNFRSF7 , is a member of the TNF-receptor superfamily limited to cells of the lymphoid lineage , and exists as both a dimeric glycoprotein on the cell surface and as a soluble protein in serum. As a type I transmembrane glycoprotein of about 55 kDa existing as disulfide-linked homodimer , CD27 has been shown to play roles in lymphoid proliferation , differentiation , and apoptosis. It has an important role in the generation of T cell immunity and is an robust marker for normal memory B cells. It is a T and B cell co-stimulatory molecule , the activity of CD27 is governed by its TNF-like ligand CD70 on lymphocytes and dendritic cells. The CD27-CD70 interaction is required for Th1 generation responses to differentiation signals and long-term maintenance of T cell immunity , and meanwhile , plays a key role in regulating B-cell differentiation , activation and immunoglobulin synthesis.

## For Research Use Only