

## PerCP/Cyanine5.5 Anti-Human CD70 Antibody[113-16]

**Catalog Number:** AN00424J

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.

### Description


<b>Reactivity</b>	Human
<b>Host</b>	Mouse
<b>Isotype</b>	Mouse IgG1, κ
<b>Clone No.</b>	113-16
<b>Isotype Control</b>	PerCP/Cyanine5.5 Mouse IgG1, κ Isotype Control[MOPC-21] [Product E-AB-F09792J]
<b>Conjugation</b>	PerCP/Cyanine 5.5
<b>Conjugation Information</b>	PerCP/Cyanine5.5 is designed to be excited by the blue laser (488 nm) and detected using an optical filter centered near 675 nm (e.g., a 690/50 nm bandpass filter).
<b>Storage Buffer</b>	Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer and 1% protein protectant.

### Applications

### Recommended usage

<b>FCM</b>	Each lot of this antibody is quality control tested by flow cytometric analysis. <b>The amount of the reagent is suggested to be used 5 μL of antibody per test (million cells in 100 μL staining volume or per 100 μL of whole blood).</b> Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.
------------	---

### Data



Staining of the U266 cell line with PerCP/Cyanine5.5 Anti-Human CD70 Antibody[113-16] (filled gray histogram) or PerCP/Cyanine5.5 Mouse IgG1, κ Isotype Control (empty black histogram). Total viable cells were used for analysis.

### Preparation & Storage

<b>Storage</b>	Keep as concentrated solution. This product can be stored at 2-8°C for 12 months. Please protected from prolonged exposure to light and do not freeze.
<b>Shipping</b>	Ice bag

### Antigen Information

<b>Alternate Names</b>	CD27 ligand;CD27L;Ki-24 antigen;Tumor necrosis factor ligand superfamily;member 7 (TNFSF7)
<b>Uniprot ID</b>	P32970

### For Research Use Only

**Gene ID**

970

**Background**

CD70, also known as CD27L, is a 50 kD type II transmembrane glycoprotein and member of the tumor necrosis factor superfamily. CD70 is expressed on activated T, B and NK cells, activated plasmacytoid dendritic cells (pDCs), and chronic B cell lymphocytic leukemia and large B cell lymphomas. CD70 costimulates T cell proliferation and differentiation. It plays a role in the pDC-induced B cell differentiation. The ligand of CD70 is CD27.

**For Research Use Only**