

Recombinant Human Artemin Protein

Catalog Number: PKSH033760

Note: *Centrifuge before opening to ensure complete recovery of vial contents.*


Description

Species	Human
Source	E.coli-derived Human Artemin protein Ala108-Gly220
Calculated MW	12.1 kDa
Observed MW	14 kDa
Accession	Q5T4W7
Bio-activity	Not validated for activity

Properties

Purity	> 95 % as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU per µg of the protein as determined by the LAL method.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from a 0.2 µm filtered solution of 20mM HEPES, 150mM NaCl, pH 7.4. Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization.
Reconstitution	Please refer to the specific buffer information in the printed manual. Please refer to the printed manual for detailed information.

Data



> 95 % as determined by reducing SDS-PAGE.

Background

For Research Use Only

Human Artemin is a GDNF family ligand that is distantly related to the TGF- β superfamily of molecules. It is synthesized as a preproprotein, and contains a variable length pre-, or signal sequence, plus a 68 amino acid (aa) proregion and a 113 aa mature segment. Following synthesis and proteolytic processing, mature ARTN is secreted as a presumably glycosylated, 28 kDa disulfide-linked homodimer that contains three intrachain disulfide bonds and the typical TGF- β signature cysteine-knot motif. In the mature region, human ARTN is 89% and 88% aa identical to rat and mouse ARTN, respectively. Human ARTN is active on rodent cells. The receptor for ARTN has been identified as the ligand binding subunit GFR α -3 plus the signal transducing subunit, RET. The GFR α -1/RET receptor complex has also been suggested to be a ligand binding unit for ARTN. ARTN is known to be a chemoattractant for sympathetic neuron axons innervating the developing cardiovascular system. It also promotes sensory neuron survival and likely plays a role in the development of the peripheral nervous system. Finally, it has been reported to reverse neuropathic pain due to nerve injury, and to help resolve morphological changes associated with nerve damage.

For Research Use Only

Toll-free: 1-888-852-8623

Web: www.elabscience.com

Tel: 1-832-243-6086

Email: techsupport@elabscience.com

Fax: 1-832-243-6017