Recombinant Human CHL-1 Protein (His Tag)

Catalog Number: PKSH031836

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description	
Species	Human
Source	HEK293 Cells-derived Human CHL-1 protein Met 1-Gln 1080, with an C-terminal His
Calculated MW	120 kDa
Observed MW	160-180 kDa
Accession	AAI04919.1
Bio-activity	Measured by the ability of the immobilized protein to support the adhesion of C6 Rat
	brain glial cells. When 5 x 10^4 cells/well are added to CHL1 coated plates (0.8 µg/ml and 100 µl/well), approximately 40%-60% will adhere specifically after 60 minutes at 37°C.
Properties	
Purity	> 95 % as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU per µg of the protein as determined by the LAL method.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80
	°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of
	reconstituted samples are stable at $< -20^{\circ}C$ for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from sterile PBS, pH 7.4
	Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants
	before lyophilization.
	Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.





> 95 % as determined by reducing SDS-PAGE.

Background

For Research Use Only

Elabscience®

Neural cell adhesion molecule L1-like protein, also known as close homolog of L1 (CHL1) is the prototypic member of the CTF / NF-1 family of transcription factors that serve as a novel calcium signaling pathway-responsive transcription factor and is considered as a member of the largest ctf complementation group, consisting of 30 of 126 ctf mutants isolated. CHL1 is a cell adhesion molecule highly related to L1. It contains structure plan of six extracellular C2-type immunoglobulin (Ig) domains followed by five fibronectin typeIII domains linked by a single membrane-spanning region to a short cytoplasmic domain. The extracellular portion of CHL1 is higyly glycosylated and involved them in hemophilic disease.