

PE/Cyanine5 Anti-Human FOLR1 Antibody[LK26]

Catalog Number: AN00483G

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description


Reactivity	Human
Host	Mouse
Isotype	Mouse IgG2a, κ
Clone No.	LK26
Isotype Control	PE/Cyanine5 Mouse IgG2a, κ Isotype Control[C1.18.4] [Product E-AB-F09802G]
Conjugation	PE/Cyanine 5
Conjugation Information	PE/Cyanine 5 is designed to be excited by the Blue (488 nm), Green (532 nm) and yellow-green (561 nm) lasers and detected using an optical filter centered near 670 nm (e.g., a 690/50 nm bandpass filter).
Storage Buffer	Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer.

Applications

Recommended usage

FCM	Each lot of this antibody is quality control tested by flow cytometric analysis. The amount of the reagent is suggested to be used 5 μL of antibody per test (million cells in 100 μL staining volume or per 100 μL of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.
------------	---

Data



Staining of the HeLa cell line with PE/Cyanine5 Anti-Human FOLR1 Antibody[LK26] (filled gray histogram) or PE/Cyanine5 Mouse IgG2a, κ Isotype Control (empty black histogram). Total viable cells were used for analysis.

Preparation & Storage

Storage	Keep as concentrated solution. This product can be stored at 2-8°C for 24 months. Please protected from prolonged exposure to light and do not freeze.
Shipping	Ice bag

Antigen Information

Alternate Names	Folate receptor alpha;FR-alpha;KB cells FBP;folate binding protein;folate receptor;adult;adult folate-binding protein;ovarian tumor-associated antigen MOV18;AN00483
Uniprot ID	P15328

For Research Use Only

Gene ID

2348

Background

Folate receptor (FOLR1), also known as folate receptor alpha, is a folate-binding protein (FBP) expressed in adults. It has high affinity for folic acid and its metabolites, including 5-methyltetrahydrofolate (5-MTF), the biologically active form of folic acid used for DNA synthesis and the cysteine cycle. FOLR1 mediates intracellular delivery of 5-MTF. FOLR1 is overexpressed in a variety of tumors, including breast, colorectal, and brain.