

A Reliable Research Partner in Life Science and Medicine

# PE Anti-Mouse CD122 Antibody[TM-Beta 1]

Catalog Number: AN00418UD

Note: Centrifuge before opening to ensure complete recovery of vial contents.

### **Description**

Reactivity Mouse Host Rat

IsotypeRat IgG2b, κClone No.TM-Beta 1

Isotype Control PE Rat IgG2b, κ Isotype Control[LTF-2] [Product E-AB-F09842D]

Conjugation PE

Conjugation Information PE is designed to be excited by the Blue (488 nm), Green (532 nm) and Yellow-Green

(561 nm) lasers and detected using an optical filter centered near 575 nm (e.g., a 585/42

nm bandpass filter).

Storage Buffer Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer.


### Applications Recommended usage

FCM Each lot of this antibody is quality control tested by flow cytometric analysis. For flow

cytometric staining, the suggested use of this reagent is  $\leq$  1.0  $\mu$ g per 10<sup>6</sup> cells in 100  $\mu$ L volume or 100  $\mu$ L of whole blood. It is recommended that the reagent be titrated for

optimal performance for each application.

#### Data



Staining of C57BL/6 murine splenocytes cells with APC Anti-Mouse NK1.1 Antibody and PE Anti-Mouse CD122 Antibody[TM-Beta 1] (left) or PE Rat IgG2b, κ Isotype Control (right). Total viable cells were used for analysis.

#### **Preparation & Storage**

**Storage** Keep as concentrated solution.

This product can be stored at 2-8°C for 12 months. Please protected from prolonged

exposure to light and do not freeze.

Shipping Ice bag

#### **Antigen Information**

Alternate Names IL-2 Receptor β chain;IL-2Rβ

 Uniprot ID
 P16297

 Gene ID
 16185

# For Research Use Only

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017

Web:www.elabscience.com Email:techsupport@elabscience.com

#### **Elabscience Bionovation Inc.**

A Reliable Research Partner in Life Science and Medicine

## **Background**

Elabscience®

CD122 is a 70-75 kD IL-2 receptor β chain also known as IL-2Rβ, which is also shared by the IL-15 receptor. It is constitutively expressed by NK cells and at lower levels by T cells, B cells, monocytes, and macrophages. The IL-2Rß chain can combine with either the common y subunit (yc, CD132) alone or with the yc subunit and the IL-2Ra subunit (CD25) to generate intermediate or high affinity IL-2 receptor complexes, respectively. CD122 expression levels can be upregulated by activation. The TM-β1 antibody does inhibit IL-2 binding to the IL-2 receptor. CD122 is expressed on murine, but not human, CD8+ Tregs involved in the maintenance of T cell homeostasis.

Fax: 1-832-243-6017 Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Web:www.elabscience.com

Rev. V1.6