Recombinant Mouse GITR/TNFRSF18 Protein (Fc & His Tag)

Catalog Number: PKSM041029

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description			
Species	Mouse		
Source	HEK293 Cells-derived Mouse GITR/TNFRSF18 protein Ser22-His153, with an C-		
	terminal Fc & His		
Calculated MW	42.3 kDa		
Observed MW	60 kDa		
Accession	O35714		
Bio-activity	Not validated for activity		
Properties			
Purity	> 90 % as determined by reducing SDS-PAGE.		
Endotoxin	< 1.0 EU per µg of the protein as determined by the LAL method.		
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80		
	°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of		
	reconstituted samples are stable at $< -20^{\circ}$ C for 3 months.		
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.		
Formulation	Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4.		
	Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants		
	before lyophilization.		
	Please refer to the specific buffer information in the printed manual.		
Reconstitution	Please refer to the printed manual for detailed information.		

Data

kDa	MK	R	
120 90 60			
40 30			
20	Sec. 1		
14	-		

> 90 % as determined by reducing SDS-PAGE.

Background

Tumor necrosis factor receptor superfamily member 18(Gitr) contains 3 TNFR-Cys repeats and it have four isforms. IsformA, isformB and isformC is single-pass type I membrane protein and isformD is a secreted protein. The protein is the receptor for TNFSF18.It seems to be involved in interactions between activated T-lymphocytes and endothelial cells and in the regulation of T-cell receptor-mediated cell death. It mediated NF-kappa-B activation via the TRAF2/NIK pathway.It binds to TRAF1, TRAF2, and TRAF3, but not TRAF5 and TRAF6 and binds through its C-terminus to SIVA 1/SIVA.It preferentially expressed in activated T lymphocytes and up-regulated in peripherical mononuclear cells after antigen stimulation/lymphocyte activation.

For Research Use Only