

Recombinant Human Contactin 3/CNTN3 Protein (708 Asp/Asn, Fc Tag)

Catalog Number: PKSH031803

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Species	Human
Source	HEK293 Cells-derived Human Contactin 3/CNTN3 protein Met 1-Ser 1002; 708 Asp/Asn, with an C-terminal hFc
Calculated MW	134.5 kDa
Observed MW	160-170 kDa
Accession	NP_065923.1
Bio-activity	Measured by the ability of the immobilized protein to support the adhesion of C6 Rat brain glial cells. When 5×10^4 cells/well are added to CNTN3 coated plates (0.8 µg/ml and 100 µl/well), approximately 30%-50% will adhere specifically after 60 minutes at 37°C.

Properties

Purity	> 90 % as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU per µg of the protein as determined by the LAL method.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from sterile 100mM Glycine, 10mM NaCl, 50mM Tris, pH 7.5 Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.

Data



> 90 % as determined by reducing SDS-PAGE.

Background

For Research Use Only

Contactins are a subgroup of molecules belonging to the immunoglobulin superfamily that are expressed exclusively in the nervous system. The subgroup consists of six members: Contactin-1; Contactin-2(TAG-1); Contactin-3(BIG-1); BIG-2; Contactin-5(NB-2) and NB-3. Since their identification in the late 1980s; Contactin-1 and Contactin-2 have been studied extensively. Axonal expression and the neurite extension activity of Contactin-1 and Contactin-2 attracted researchers to study the function of these molecules in axon guidance during development. Contactin-1 and Contactin-2 have come to be known as the principal molecules in the function and maintenance of myelinated neurons. In contrast; the function of the other four members of this subgroup remained unknown until recently. Contactin-3; also known as CNTN3 (BIG-1 in rat and PANG in mouse); is a GPI-linked glycoprotein that is expressed on cerebellar Purkinje cells; amygdaloid and thalamic neurons and olfactory granule cells. In the brain; Contactin-3 is expressed in frontal lobe; occipital lobe; cerebellum and amygdala. Contactin-3 contains 4 fibronectin type-III domains and 6 Ig-like C2-type (immunoglobulin-like) domains. Human Contactin-3 shares 92% aa identity with mouse Contactin-3. The exact function of Contactin-3 is unclear. Contactin-3 may mediate cell-cell interaction and may promote neurite outgrowth.

For Research Use Only

Toll-free: 1-888-852-8623
Web: www.elabscience.com

Tel: 1-832-243-6086
Email: techsupport@elabscience.com

Fax: 1-832-243-6017