

Recombinant CD50/ICAM3 Monoclonal Antibody

catalog number: AN301481L

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Reactivity	Human;
Immunogen	Recombinant human CD50/ICAM3 fragment
Host	Rabbit
Isotype	IgG, κ
Clone	A176
Purification	Protein A purified
Buffer	PBS, 50% glycerol, 0.05% Proclin 300, 0.05% protein protectant.


Applications Recommended Dilution

IHC	1:100-1:500
IF	1:100-1:200
FCM	1:50-1:100


Data




Immunohistochemistry of paraffin-embedded Human colon using CD50/ICAM3 Monoclonal Antibody at dilution of 1:500.




Immunohistochemistry of paraffin-embedded Human tonsil using CD50/ICAM3 Monoclonal Antibody at dilution of 1:1000.



Immunofluorescent analysis of (4% Paraformaldehyde) fixed Ramos cells using anti-CD50/ICAM3 Monoclonal Antibody at dilution of 1:200.



Immunofluorescent analysis of (4% Paraformaldehyde) fixed Jurkat cells using anti-CD50/ICAM3 Monoclonal Antibody at dilution of 1:200.



For Research Use Only

Toll-free: 1-888-852-8623
Web: www.elabscience.com

Tel: 1-832-243-6086
Email: techsupport@elabscience.com

Fax: 1-832-243-6017

Rev. V1.0

Flow cytometric analysis of human CD50/ICAM3 expression on PBMC cells. Cells were stained with purified anti-Human CD50/ICAM3, then a Alexa Fluor 488-conjugated second step antibody. The histogram were derived from events with the forward and side light-scatter characteristics of intact cells.

Preparation & Storage

Storage	Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.
Shipping	Ice bag

Background

ICAM proteins are ligands for the leukocyte adhesion protein LFA-1 (integrin alpha-L/beta-2). ICAM3 is also a ligand for integrin alpha-D/beta-2. In association with integrin alpha-L/beta-2, contributes to apoptotic neutrophil phagocytosis by macrophages.

For Research Use Only