



A Reliable Research Partner in Life Science and Medicine

## Elab Fluor® 488 Anti-Mouse CD107a/LAMP-1 Antibody[1D4B]

Catalog Number: E-AB-F1254L

Note: Centrifuge before opening to ensure complete recovery of vial contents.

**Description** 

Reactivity Mouse Rat Host

Isotype Rat IgG2a, ĸ 1D4B Clone No.

Isotype Control Elab Fluor® 488 Rat IgG2a, κ Isotype Control[2A3] [Product E-AB-F09832L]

Conjugation Elab Fluor®488

**Conjugation Information** Elab Fluor® 488 is designed to be excited by the Blue laser (488 nm) and detected using

an optical filter centered near 520 nm (e.g., a 525/40 nm bandpass filter).

Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer. Storage Buffer

**Applications** Recommended usage

Each lot of this antibody is quality control tested by flow cytometric analysis. The amount **FCM** 

> of the reagent is suggested to be used 5 µL of antibody per test (million cells in 100 μL staining volume or per 100 μL of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for

individual use.

**Preparation & Storage** 

Storage Keep as concentrated solution.

This product can be stored at 2-8°C for 12 months. Please protected from prolonged

exposure to light and do not freeze.

**Shipping** Ice bag

**Antigen Information** 

**Alternate Names** CD107 antigen-like family member Alysosomal membrane glycoprotein 1lysosome-

associated membrane protein 1;LAMP-1;Lysosome-associated membrane

Associated Membrane Protein 1 (LAMP-1). CD107a is one of the two major

glycoprotein 1;LAMP-1

**Uniprot ID** P11438 Gene ID 16783

**Background** The 1D4B antibody recognizes CD107a which is also known as, Lysosome-

> glycoproteins in lysosome membranes that provide useful markers to distinguish lysosomes from other organelles. CD107a may play a role in the lysosomal degradation of certain molecules. Mouse CD107a is a type I transmembrane glycoprotein. It consists of a 40-kDa core protein which is heavily glycosylated to form heterogeneous mature glycoprotein of 110-140 kDa. It is principally expressed in epithelial cells and macrophages in a variety of organs. Following activation, CD107a is relocated to the surface of some lymphocytes, macrophages, epithelial cells, endothelial cells, platelets, and tumor cells. Cell-surface CD107a may participate in intercellular adhesion and adhesion to the extracellular matrix. Cell surface CD107a expression can serve as a useful marker for cytotoxic NK and CD8+ T cells, as well as,

> > Rev. V1.8

some malignant tumor cells.

## For Research Use Only

Tel: 1-832-243-6086 Fax: 1-832-243-6017 Toll-free: 1-888-852-8623 Web:www.elabscience.com