

Mouse OxLDL Antibody Pair Set

Catalog No.	E-KAB-0682	Applications	ELISA
Synonyms	CLEC8A;CLEC8ASLOX1;C-type lectin domain family 8 member A;hLOX-1;Lectin-like oxidized LDL receptor 1;Lectin-like oxLDL receptor 1;Lectin-type oxidized LDL receptor 1;LOX1;LOX-1;LOX1ox LDL receptor 1;LOXIN;OLR1		

Kit components & Storage

Title	Specifications	Storage
Mouse OxLDL Capture Antibody	1 vial, 100 µg	Store at -20°C. Avoid freeze / thaw cycles.
Mouse OxLDL Detection Antibody (Biotin)	1 vial, 50 µL	Store at -20°C. Avoid freeze / thaw cycles.

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Product Information

Items		Characteristic (E-KAB-0682)	
		Mouse OxLDL Capture Antibody	Mouse OxLDL Detection Antibody (Biotin)
Immunogen Information	Immunogen	Recombinant Mouse OxLDL protein	Recombinant Mouse OxLDL protein
	Swissprot	/	
Product details	Reactivity	Mouse	Mouse
	Host	Goat	Goat
	Conjugation	Unconjugated	Biotin
	Concentration	0.5mg/mL	/
	Buffer	PBS with 0.04% Proclin 300, 50% glycerol, pH 7.4	PBS with 0.04% Proclin 300, 1% protective protein, 50% glycerol, pH 7.4
	Purify	Affinity purification	Affinity purification
	Specificity	Detects Mouse OxLDL in ELISAs.	

For Research Use Only

Applications

Mouse OxLDL Sandwich ELISA Assay:

	Recommended Concentration/Dilution	Reagent	Images										
ELISA Capture	0.5-4ug/mL	Mouse OxLDL Capture Antibody	 <table border="1"> <caption>Standard Curve Data (Approximate)</caption> <thead> <tr> <th>Mouse OxLDL concentration (ng/mL)</th> <th>Optical Density</th> </tr> </thead> <tbody> <tr><td>0.1</td><td>0.15</td></tr> <tr><td>1</td><td>0.25</td></tr> <tr><td>10</td><td>0.5</td></tr> <tr><td>100</td><td>1.5</td></tr> </tbody> </table>	Mouse OxLDL concentration (ng/mL)	Optical Density	0.1	0.15	1	0.25	10	0.5	100	1.5
Mouse OxLDL concentration (ng/mL)	Optical Density												
0.1	0.15												
1	0.25												
10	0.5												
100	1.5												
ELISA Detection	1:1000-1:10000	Mouse OxLDL Detection Antibody (Biotin)											

Note: This standard curve is only for demonstration purposes. A standard curve should be generated for each assay!

Background

Receptor that mediates the recognition, internalization and degradation of oxidatively modified low density lipoprotein (oxLDL) by vascular endothelial cells. OxLDL is a marker of atherosclerosis that induces vascular endothelial cell activation and dysfunction, resulting in pro-inflammatory responses, pro-oxidative conditions and apoptosis. Its association with oxLDL induces the activation of NF-kappa-B through an increased production of intracellular reactive oxygen and a variety of pro-atherogenic cellular responses including a reduction of nitric oxide (NO) release, monocyte adhesion and apoptosis. In addition to binding oxLDL, it acts as a receptor for the HSP70 protein involved in antigen cross-presentation to naive T-cells in dendritic cells, thereby participating in cell-mediated antigen cross-presentation. Also involved in inflammatory process, by acting as a leukocyte-adhesion molecule at the vascular interface in endotoxin-induced inflammation. Also acts as a receptor for advanced glycation end (AGE) products, activated platelets, monocytes, apoptotic cells and both Gram-negative and Gram-positive bacteria.