

## Recombinant Mouse PLA2G7/Lp-PLA2 Protein (His Tag)

**Catalog Number:** PKSM040287

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.


### Description

<b>Species</b>	Mouse
<b>Source</b>	HEK293 Cells-derived Mouse PLA2G7/Lp-PLA2 protein Met1-Asn440, with an C-terminal His
<b>Calculated MW</b>	48.3 kDa
<b>Accession</b>	NP_038765.2
<b>Bio-activity</b>	Not validated for activity

### Properties

<b>Purity</b>	> 90 % as determined by reducing SDS-PAGE.
<b>Endotoxin</b>	< 1.0 EU per µg of the protein as determined by the LAL method.
<b>Storage</b>	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80 °C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
<b>Shipping</b>	This product is provided as lyophilized powder which is shipped with ice packs.
<b>Formulation</b>	Lyophilized from sterile PBS, pH 7.4 Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
<b>Reconstitution</b>	Please refer to the printed manual for detailed information.

### Data



> 90 % as determined by reducing SDS-PAGE.

### Background

### For Research Use Only

Platelet-activating factor acetylhydrolase, also known as 1-alkyl-2-acetyl-glycerophosphocholine esterase, 2-acetyl-1-alkylglycero-phosphocholine esterase, Group-VIIA phospholipase A2, LDL-associated phospholipase A2, PAF 2-acetylhydrolase, PLA2G7 and PAFAH, is a secreted protein which belongs to the AB hydrolase superfamily and Lipase family. PLA2G7 / PAFAH modulates the action of platelet-activating factor (PAF) by hydrolyzing the sn-2 ester bond to yield the biologically inactive lyso-PAF. It has a specificity for substrates with a short residue at the sn-2 position. It is inactive against long-chain phospholipids. PLA2G7 / PAFAH is a potent pro- and anti-inflammatory molecule that has been implicated in multiple inflammatory disease processes, including cardiovascular disease. PLA2G7 also represents an important, potentially functional candidate in the pathophysiology of coronary artery disease (CAD). Defects in PLA2G7 are the cause of platelet-activating factor acetylhydrolase deficiency (PLA2G7 deficiency). It is a trait which is present in 27% of Japanese. It could have a significant physiologic effect in the presence of inflammatory bodily responses.