## **Elabscience**®


## Elab Fluor<sup>®</sup> Violet 610 Anti-Mouse CD19 Antibody[1D3]

Catalog Number: E-AB-F0986UT

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description	
Reactivity	Mouse
Host	Rat
Isotype	Rat lgG2a, к
Clone No.	1D3
Isotype Control	Elab Fluor <sup>®</sup> Violet 610 Rat IgG2a, к Isotype Control[2A3] [Product E-AB-F09833T]
Conjugation	Elab Fluor <sup>®</sup> Violet 610
Conjugation Information	Elab Fluor <sup>®</sup> Violet 610 is designed to be excited by the violet laser (405 nm) and detected using an optical filter centered near 613 nm (e.g., a 615/20 nm bandpass filter).
Storage Buffer	Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer and 1% protein protectant.
Applications	Recommended usage
FCM	Each lot of this antibody is quality control tested by flow cytometric analysis. Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use. We suggest each investigator should titrate the reagent to obtain optimal results [The recommended concentration is 0.1-1 µg/10 <sup>6</sup> cells

reagent to obtain optimal results [The recommended concentration is 0.1-1  $\mu$ g/10<sup>6</sup> cells in 100  $\mu$ L volume].



Staining of C57BL/6 murine splenocytes with FITC Anti-

Mouse CD3 Antibody[17A2] and Elab Fluor<sup>®</sup> Violet 610 Anti-Mouse CD19 Antibody[1D3] (left) or Elab Fluor<sup>®</sup> Violet 610 Rat IgG2a,  $\kappa$  Isotype Control (right). Total viable cells were used for analysis.

Keep as concentrated solution.
This product can be stored at 2-8°C for 12 months. Please protected from prolonged exposure to light and do not freeze.
lce bag
B-lymphocyte antigen CD19;CD19;Cd19;Differentiation antigen CD19
P25918

## For Research Use Only

Toll-free: 1-888-852-8623 Web:<u>w w w .elabscience.com</u>

## **Elabscience**®

Gene ID Background 12478

CD19 is a 95 kD glycoprotein also known as B4. It is a member of the lg superfamily, expressed on all pro-B to mature B cells (during development) and follicular dendritic cells. Plasma cells do not express CD19. CD19, in association with CD21 and CD81, forms a molecular complex integral to B cell activation.