# IGFBP-6 Polyclonal Antibody(Capture/Detector)

catalog number: AN002460P



Note: Centrifuge before opening to ensure complete recovery of vial contents.

| -            |   |   |    |    |    | ٠, |    |   |
|--------------|---|---|----|----|----|----|----|---|
| D            | Α |   | C  | PТ | m  | ŤΤ | U. | m |
| $\mathbf{L}$ | v | 0 | v. | ш  | עו | UΙ | U. |   |

**Reactivity** Mouse

Immunogen Recombinant Mouse IGFBP-6 protein expressed by Mammalian

Host Rabbit
Isotype Rabbit IgG

**Purification** Antigen Affinity Purification


**Conjugation** Unconjugated

**buffer** Phosphate buffered solution, pH 7.2, containing 0.05% proclin 300.

**Applications** Recommended Dilution

**ELISA Capture** 2-8 μg/mL **ELISA Detector** 0.1-0.4 μg/mL

#### Data



Sandwich ELISA-Recombinant Mouse IGFBP-6 protein standard curve. Background subtracted standard curve using IGFBP-6 antibody(AN002460P)(Capture),IGFBP-6 antibody(AN002460P)(Detector) in sandwich ELISA. The reference range value for Recombinant Mouse IGFBP-6 protein is 7.81-500 pg/mL.

### **Preparation & Storage**

Storage Storage Store at 4°C valid for 12 months or -20°C valid for long term storage, avoid freeze /

thaw cycles.

**Shipping** The product is shipped with ice pack, upon receipt, store it immediately at the

temperature recommended.

## Background

The superfamily of insulin-like growth factor (IGF) binding proteins include the six high-affinity IGF binding proteins (IGFBP) and at least four additional low-affinity binding proteins referred to as IGFBP related proteins (IGFBP-rP). All IGFBP superfamily members are cysteine-rich proteins with conserved cysteine residues, which are clustered in the amino- and carboxy-terminal thirds of the molecule. IGFBPs modulate the biological activities of IGF proteins. Some IGFBPs may also have intrinsic bioactivity that is independent of their ability to bind IGF proteins. Post-translational modifications of IGFBPs, including glycosylation, phosphorylation and proteolysis, have been shown to modify the affinities of the binding proteins to IGF.

## For Research Use Only