A Reliable Research Partner in Life Science and Medicine

Recombinant Mouse VEGF-A/VEGF164 Protein

Catalog Number: PKSM041180

Note: Centrifuge before opening to ensure complete recovery of vial contents.

\mathbf{r}					٠.			
H))	es	C	r٦	n	tı	n	m	

Species Mouse

Source P.pastoris-derived Mouse VEGF-A/VEGF164 protein Ala27-Arg190

 Mol_Mass
 19.3 kDa

 Accession
 Q00731-2

Bio-activity Not validated for activity

Properties

Purity > 90 % as determined by reducing SDS-PAGE.

Endotoxin < 1.0 EU per µg of the protein as determined by the LAL method.

Storage Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80

°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of

reconstituted samples are stable at < -20°C for 3 months.

ShippingThis product is provided as lyophilized powder which is shipped with ice packs.FormulationLyophilized from a 0.2 μm filtered solution of 20mM PB, 250mM NaCl, pH7.0.


Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants

before lyophilization.

Please refer to the specific buffer information in the printed manual.

Reconstitution Please refer to the printed manual for detailed information.

Data



Background

Mouse Vascular endothelial growth factor (VEGF or VEGF-A), is a potent mediator of both angiogenesis and vasculogenesis in the fetus and adult. It is a member of the PDGF/VEGF growth factor family that is characterized by a cystine knot structure formed by eight conserved cysteine residues. Alternately spliced isoforms of 120, 164 and 188 aa found in mouse. VEGF binds the type I transmembrane receptor tyrosine kinases VEGF R1 (also called Flt-1) and VEGF R2 (Flk-/KDR) on endothelial cells. Although affinity is highest for binding to VEGF R1, VEGF R2 appears to be the primary mediator of VEGF angiogenic activity. VEGF is required during embryogenesis to regulate the proliferation, migration, and survival of endothelial cells. It may play a role in increasing vascular permeability during lactation, when increased transport of molecules from the blood is required for efficient milk protein synthesis.

For Research Use Only