

Recombinant Human Sonic Hedgehog/SHH Protein (aa 1-197, His Tag)

Catalog Number: PKSH031645

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Species	Human
Source	HEK293 Cells-derived Human Sonic Hedgehog/SHH protein Met 1-Gly 197, with an C-terminal His
Calculated MW	21 kDa
Observed MW	24 kDa
Accession	Q15465
Bio-activity	Measured by its ability to induce alkaline phosphatase production by C3H10T1/2 mouse embryonic fibroblast cells. The ED ₅₀ for this effect is typically 2-10 µg/mL.

Properties

Purity	> 95 % as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU per µg of the protein as determined by the LAL method.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from sterile PBS, pH 7.4 Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.

Data



Measured by its ability to induce alkaline phosphatase production by C3H10T1/2 mouse embryonic fibroblast cells.

Nakamura, T. et al. (1997) Biochem. Biophys. Res. Commun. 237:465. The ED₅₀ for this effect is typically 2-10 µg/mL.

Background

For Research Use Only

Toll-free: 1-888-852-8623
Web: www.elabscience.com

Tel: 1-832-243-6086
Email: techsupport@elabscience.com

Fax: 1-832-243-6017

Sonic HedgeHog; also known as sonic hedgehog protein; belongs to the hedgehog family. It cannot be detected in adult tissues while can be found in fetal intestine; liver; lung; and kidney. Sonic HedgeHog is a protein that is vital in guiding the early embryo. It has been associated as the major inductive signal in patterning of the ventral neural tube; the anterior-posterior limb axis; and the ventral somites. Sonic HedgeHog intercellular signal is essential for a various patterning events during development: signal produced by the notochord that induces ventral cell fate in the neural tube and somites; and the polarizing signal for patterning of the anterior-posterior axis of the developing limb bud. Sonic HedgeHog binds to the patched receptor; which functions in association with smoothened; to activate the transcription of target genes. In the absence of sonic HedgeHog; patched receptor represses the constitutive signaling activity of smoothened. Sonic HedgeHog also regulates another factor; the gli oncogene. Defects in sonic hedgehog can cause microphthalmia isolated with coloboma type 5; triphalangeal thumb-polysyndactyly syndrome and holoprosencephaly type 3.

For Research Use Only

Toll-free: 1-888-852-8623
Web: www.elabscience.com

Tel: 1-832-243-6086
Email: techsupport@elabscience.com

Fax: 1-832-243-6017