

## Human RET Antibody Pair Set

<b>Catalog No.</b>	E-KAB-0239	<b>Applications</b>	ELISA
<b>Synonyms</b>	Proto-oncogene tyrosine-protein kinase receptor Ret, Cadherin family member 12, Proto-oncogene c-Ret		

### Kit components & Storage

Title	Specifications	Storage
Human RET Capture Antibody	1 vial, 100 µg	Store at -20℃ for one year. Avoid freeze / thaw cycles.
Human RET Detection Antibody (Biotin)	1 vial, 50 µL	Store at -20℃ for one year. Avoid freeze / thaw cycles.

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.


### Product Information

Items		Characteristic (E-KAB-0239)	
		Human RET Capture Antibody	Human RET Detection Antibody (Biotin)
Immunogen Information	Immunogen	Recombinant Human RET protein	Recombinant Human RET protein
	Swissprot	P07949	
Product details	Reactivity	Human	Human
	Host	Goat	Goat
	Conjugation	Unconjugated	Biotin
	Concentration	0.5mg/mL	/
	Buffer	PBS with 0.04% Proclin 300, 50% glycerol, pH 7.4	PBS with 0.04% Proclin 300, 1% protective protein, 50% glycerol, pH 7.4
	Purify	Antigen Affinity	Antigen Affinity
	Specificity	Detects Human RET in ELISAs.	

### For Research Use Only

## Applications

### Human RET Sandwich ELISA Assay:

	Recommended Concentration/Dilution	Reagent	Images
ELISA Capture	0.5-4µg/mL	Human RET Capture Antibody	
ELISA Detection	1:1000-1:10000	Human RET Detection Antibody (Biotin)	

**Note:** This standard curve is only for demonstration purposes. A standard curve should be generated for each assay!

## Background

This gene, a member of the cadherin superfamily, encodes one of the receptor tyrosine kinases, which are cell-surface molecules that transduce signals for cell growth and differentiation. This gene plays a crucial role in neural crest development, and it can undergo oncogenic activation in vivo and in vitro by cytogenetic rearrangement. Mutations in this gene are associated with the disorders multiple endocrine neoplasia, type IIA, multiple endocrine neoplasia, type IIB, Hirschsprung disease, and medullary thyroid carcinoma. Two transcript variants encoding different isoforms have been found for this gene. Additional transcript variants have been described but their biological validity has not been confirmed. RET (Ret Proto-Oncogene) is a Protein Coding gene. Diseases associated with RET include Multiple Endocrine Neoplasia Iia and Medullary Thyroid Carcinoma, Familial. Among its related pathways are RET signaling and Innate Immune System. GO annotations related to this gene include calcium ion binding and protein kinase activity. An important paralog of this gene is PDGFRB.

## For Research Use Only