Elabscience Biotechnology Co., Ltd.



A Reliable Research Partner in Life Science and Medicine

MMP2 Polyclonal Antibody

catalog number: E-AB-32054

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Reactivity Human; Mouse; Rat; Monkey

Immunogen Synthesized peptide derived from the C-terminal region of human MMP-2

Host Rabbit Isotype IgG

PurificationAffinity purificationConjugationUnconjugated

Buffer Phosphate buffered solution, pH 7.4, containing 0.05% stabilizer, 0.5% protein

protectant and 50% glycerol.

Applications	Recommended Dilution
WB	1:500-1:2000
IHC	1:100-1:300
IF	1:200-1:1000

Data

(kD) HT29


170130957255-

HT-29 cells using MMP2

Immunohistochemistry of paraffin embedder

Western Blot analysis of HT-29 cells using MMP2 Polyclonal Antibody at dilution of 1:1000.

Observed-MW:74 kDa Calculated-MW:74 kDa



Immunohistochemistry of paraffin-embedded Human liver cancer tissue using MMP2 Polyclonal Antibody at dilution of 1:200.

Immunofluorescence analysis of Rat lung tissue using MMP2 Polyclonal Antibody at dilution of 1:200.

Preparation & Storage

Storage Storage Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.

Shipping The product is shipped with ice pack, upon receipt, store it immediately at the

Web: www.elabscience.cn

temperature recommended.

Background

For Research Use Only

Tel: 400-999-2100

Elabscience®

Elabscience Biotechnology Co., Ltd.

A Reliable Research Partner in Life Science and Medicine

Ubiquitinous metalloproteinase that is involved in diverse functions such as remodeling of the vasculature, angiogenesis, tissue repair, tumor invasion, inflammation, and atherosclerotic plaque rupture. As well as degrading extracellular matrix proteins, can also act on several nonmatrix proteins such as big endothelial 1 and beta-type CGRP promoting vasoconstriction. Also cleaves KISS at a Gly-Leu bond. Appears to have a role in myocardial cell death pathways. Contributes to myocardial oxidative stress by regulating the activity of GSK3beta. Cleaves GSK3beta in vitro. PEX, the C-terminal non-catalytic fragment of MMP2, posseses anti-angiogenic and anti-tumor properties and inhibits cell migration and cell adhesion to FGF2 and vitronectin. Ligand for integrinv/beta3 on the surface of blood vessels.

For Research Use Only

Tel: 400-999-2100 Web: www.elabscience.cn Email:techsupport@elabscience.cn