

A Reliable Research Partner in Life Science and Medicine

APC Anti-Rat CD4(domain 1) Antibody[OX-38]

Catalog Number: E-AB-F1105E

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Reactivity Rat Mouse Host

Isotype Mouse IgG2a, ĸ

Clone No. OX-38

APC Mouse IgG2a, κ Isotype Control[C1.18.4] [Product E-AB-F09802E] Isotype Control

APC Conjugation

Conjugation Information APC is designed to be excited by the Red (627-640 nm) laser and detected using an

optical filter centered near 660 nm (e.g., a 660/20 nm bandpass filter).


Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer. Storage Buffer

Applications Recommended usage

FCM

Each lot of this antibody is quality control tested by flow cytometric analysis. The amount of the reagent is suggested to be used 5 µL of antibody per test (million cells in 100 μL staining volume or per 100 μL of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.

Data



Rat splenocytes are stained with APC Anti-Rat CD4(domain 1) Antibody (filled gray histogram). Unstained splenocytes (empty black histogram) are used as control.

Preparation & Storage

Keep as concentrated solution. Storage

This product can be stored at 2-8°C for 12 months. Please protected from prolonged

exposure to light and do not freeze.

Shipping Ice bag

Antigen Information

Alternate Names CD4;T-cell surface antigen T4/Leu-3;T-cell surface glycoprotein CD4;domain 1

Uniprot ID P05540 Gene ID 24932

For Research Use Only

Fax: 1-832-243-6017 Tel: 1-832-243-6086 Toll-free: 1-888-852-8623 Web:www.elabscience.com

Rev. V1.5

Elabscience Bionovation Inc.



A Reliable Research Partner in Life Science and Medicine

Background

CD4, also known as T4, is a 55kD glycoprotein member of the immunoglobin superfamily and is expressed on majority of thymocytes, macrophages, and a peripheral T cell subset (T helper cells). CD4 is a T cell co-receptor that interacts with the MHC class II molecule and is involved in T cell activation.