

GNAT1 Polyclonal Antibody

catalog number: E-AB-19919

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description


Reactivity	Human;Mouse
Immunogen	Synthetic peptide of human GNAT1
Host	Rabbit
Isotype	IgG
Purification	Antigen affinity purification
Conjugation	Unconjugated
buffer	Phosphate buffered solution, pH 7.4, containing 0.05% stabilizer and 50% glycerol.

Applications

Recommended Dilution


WB	1:500-1:2000
IHC	1:30-1:150

Data



Western blot analysis of Human fetal liver tissue lysate using GNAT1 Polyclonal Antibody at dilution of 1:250

Observed-MV:Refer to figures
Calculated-MV:40 kDa



Immunohistochemistry of paraffin-embedded Human colorectal cancer tissue using GNAT1 Polyclonal Antibody at dilution of 1:25(×200)

Preparation & Storage

Storage	Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.
Shipping	The product is shipped with ice pack,upon receipt,store it immediately at the temperature recommended.

Background

Transducin is a 3-subunit guanine nucleotide-binding protein (G protein) which stimulates the coupling of rhodopsin and cGMP-phosphodiesterase during visual impulses. The transducin alpha subunits in rods and cones are encoded by separate genes. This gene encodes the alpha subunit in rods. This gene is also expressed in other cells, and has been implicated in bitter taste transduction in rat taste cells. Mutations in this gene result in autosomal dominant congenital stationary night blindness. Multiple alternatively spliced variants, encoding the same protein, have been identified. GNAT1 (GProtein Subunit Alpha Transducin 1) is a Protein Coding gene. Diseases associated with GNAT1 include Night Blindness, Congenital Stationary, Autosomal Dominant 3 and Night Blindness, Congenital Stationary, Type 1G. Among its related pathways are Phospholipase-C Pathway and Phototransduction. GO annotations related to this gene include GTP binding and GTPase activity. An important paralog of this gene is GNAT2.

For Research Use Only