

Recombinant Human XIAP/BIRC4 Protein (AVI Tag)

Catalog Number:PKSH031510



Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Synonyms	API3;BIRC4;hIAP-3;hIAP3;IAP-3;ILP1;MIHA;XLP2
Species	Human
Expression Host	E.coli
Sequence	Leu 121-Thr 356
Accession	NP_001158.2
Calculated Molecular Weight	29.1 kDa
Observed molecular weight	29.1 kDa
Tag	C-Avi
Bioactivity	Measured by its binding ability in a functional ELISA. Immobilized recombinant human SMAC-His at 10 µg/ml (100 µl/well) can bind recombinant human XIAP-AVI with a linear range of 0.125-1.0 µg/ml.

Properties

Purity	> 75 % as determined by reducing SDS-PAGE.
Endotoxin	Please contact us for more information.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from sterile 25mM Tris, 10mM DTT, 1% glycerol, 0.2M Glutamine Potassium, pH 8.0 Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in
Reconstitution	Please refer to the printed manual for detailed information.

Data



> 75 % as determined by reducing SDS-PAGE.

Background

E3 ubiquitin-protein ligase XIAP / BIRC4, also known as inhibitor of apoptosis protein 3, X-linked inhibitor of apoptosis protein, and IAP-like protein, is a protein that belongs to a family of apoptotic suppressor proteins. Members of this family share a conserved motif termed, baculovirus IAP repeat, which is necessary for their anti-apoptotic function. XIAP / BIRC4 functions through binding to tumor necrosis factor receptor-associated factors TRAF1 and TRAF2 and

For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623

Web: www.elabscience.com

Tel: 1-832-243-6086

Email: techsupport@elabscience.com

Fax: 1-832-243-6017

Recombinant Human XIAP/BIRC4 Protein (AVI Tag)

Catalog Number:PKSH031510



inhibits apoptosis induced by menadione, a potent inducer of free radicals, and interleukin 1-beta converting enzyme. XIAP / BIRC4 also inhibits at least two members of the caspase family of cell-death proteases, caspase-3 and caspase-7. Mutations in this encoding gene are the cause of X-linked lymphoproliferative syndrome. Alternate splicing results in multiple transcript variants. Thought to be the most potent apoptosis suppressor, XIAP / BIRC4, directly binds and inhibits caspases -3, -7 and -9. Survivin, which also binds to several caspases, is up-regulated in a many tumour cell types. Defects in XIAP / BIRC4 are the cause of lymphoproliferative syndrome X-linked type 2 (XLP2). XLP is a rare immunodeficiency characterized by extreme susceptibility to infection with Epstein-Barr virus (EBV). Symptoms include severe or fatal mononucleosis, acquired hypogammaglobulinemia, pancytopenia and malignant lymphoma.

For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623

Web: www.elabscience.com

Tel: 1-832-243-6086

Email: techsupport@elabscience.com

Fax: 1-832-243-6017