A Reliable Research Partner in Life Science and Medicine

## Recombinant Mouse SHP2/PTPN11 Protein (His Tag)

Catalog Number: PKSM040671

Note: Centrifuge before opening to ensure complete recovery of vial contents.

#### Description

**Species** Mouse

Source HEK293 Cells-derived Mouse SHP2/PTPN11 protein Met1-Arg593, with an C-terminal

His

Calculated MW 69.5 kDa
Observed MW 65 kDa
Accession P35235-2

**Bio-activity** Measured by its ability to dephosphorylate a tyrosine residue in a peptide containing

the EGFR Y992 phosphorylation site (Catalog # ES006). The specific activity is > 1

pmoles/min/µg.

#### **Properties**

**Purity** > 90 % as determined by reducing SDS-PAGE.

Endotoxin < 1.0 EU per µg of the protein as determined by the LAL method.

Storage Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80

°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of

reconstituted samples are stable at < -20°C for 3 months.

**Shipping** This product is provided as lyophilized powder which is shipped with ice packs.

**Formulation** Lyophilized from sterile PBS, pH 7.4


Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants

before lyophilization.

Please refer to the specific buffer information in the printed manual.

**Reconstitution** Please refer to the printed manual for detailed information.

#### Data



> 90 % as determined by reducing SDS-PAGE.

### Background

# Elabscience®

#### **Elabscience Bionovation Inc.**

A Reliable Research Partner in Life Science and Medicine

SHP2, also known as PTPN11, belongs to the protein-tyrosine phosphatase(PTP) family, non-receptor class 2 subfamily. PTPs catalyze the removal of phosphate groups from tyrosine residues by the hydrolysis of phosphoric acid monoester s. They dephosphorylate EGFR, JAK2 and TYK2 kinases, promoting oncogenic transformation. SHP2 is widely expressed, with highest levels in heart, brain, and skeletal muscle. SHP2 acts downstream of various receptor and cytoplasmic protein tyrosine kinases to participate in the signal transduction from the cell surface to the nucleus. It also dephosphorylates ROCK2 at Tyr-722 resulting in stimulatation of its RhoA binding activity.

Toll-free: 1-888-852-8623 Web:www.elabscience.com Tel: 1-832-243-6086 Email:techsupport@elabscience.com Fax: 1-832-243-6017