

## Recombinant Human PDGF-AA Protein (His Tag)

**Catalog Number:** PKSH032905

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.


### Description

<b>Species</b>	Human
<b>Source</b>	E.coli-derived Human PDGF-AA protein Ser87-Thr 211, with an N-terminal His
<b>Calculated MW</b>	15.9 kDa
<b>Observed MW</b>	16 kDa
<b>Accession</b>	P04085
<b>Bio-activity</b>	Not validated for activity

### Properties

<b>Purity</b>	> 95 % as determined by reducing SDS-PAGE.
<b>Endotoxin</b>	< 1.0 EU per µg of the protein as determined by the LAL method.
<b>Storage</b>	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80 °C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
<b>Shipping</b>	This product is provided as lyophilized powder which is shipped with ice packs.
<b>Formulation</b>	Lyophilized from a 0.2 µm filtered solution of 4mM HCl. Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
<b>Reconstitution</b>	Please refer to the printed manual for detailed information.

### Data



> 95 % as determined by reducing SDS-PAGE.

### Background

Platelet-derived growth factor subunit A (PDGFA); belongs to the PDGF/VEGF growth factor family. PDGFA is a secreted protein; stored in platelet alpha-granules and released by platelets upon wounding. PDGFA is potent mitogens for a variety of cell types including smooth muscle cells; connective tissue cells; bone and cartilage cells; and some blood cells. It plays an essential role in the regulation of embryonic development; cell proliferation; cell migration; survival and chemotaxis. PDGFA is required for normal lung alveolar septum formation during embryogenesis; normal development of the gastrointestinal tract; normal development of Leydig cells and spermatogenesis; normal oligodendrocyte development and normal myelination in the spinal cord and cerebellum. It plays an important role in wound healing; Signaling is modulated by the formation of heterodimers with PDGFB.

### For Research Use Only

Toll-free: 1-888-852-8623  
Web: [www.elabscience.com](http://www.elabscience.com)

Tel: 1-832-243-6086  
Email: [techsupport@elabscience.com](mailto:techsupport@elabscience.com)

Fax: 1-832-243-6017