# **Elabscience**®

# LIG3 Polyclonal Antibody

### catalog number: E-AB-64684

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description	
Reactivity	Human;Mouse;Rat
Immunogen	Recombinant fusion protein of human LIG3 (NP_039269.2).
Host	Rabbit
Isotype	IgG
Purification	Affinity purification
Buffer	Phosphate buffered solution, pH 7.4, containing 0.05% stabilizer and 50% glycerol.
Applications	Recommended Dilution

Applications	Recommended Dilution
WB	1:500-1:2000
IHC	1:50-1:200

#### Data



Western blot analysis of extracts of various cell lines using Immunohistochemistry of paraffin-embedded Rat heart using

LIG3 Polyclonal Antibody at dilution of 1:1000.

### Observed-MW:112 kDa

#### Calculated-MW:95 kDa/102 kDa/106 kDa/112 kDa



Immunohistochemistry of paraffin-embedded Human oophoroma using LIG3 Polyclonal Antibody at dilution of



Immunohistochemistry of paraffin-embedded Mouse heart using LIG3 Polyclonal Antibody at dilution of 1:100 (40x lens).

## For Research Use Only

Toll-free: 1-888-852-8623 Web:<u>w w .elabscience.com</u>



LIG3 Polyclonal Antibody at dilution of 1:100 (40x lens).



Immunohistochemistry of paraffin-embedded Human esophageal cancer using LIG3 Polyclonal Antibody at dilution of 1:100 (40x lens).

# **Elabscience**®

Elabscience Bionovation Inc. A Reliable Research Partner in Life Science and Medicine

Storage Store at -20	°C Valid for 12 months. Avoid freeze / thaw cycles.
Shipping The product temperature	t is shipped with ice pack, upon receipt, store it immediately at the recommended.

#### Background

This gene is a member of the DNA ligase family. Each member of this family encodes a protein that catalyzes the joining of DNA ends but they each have a distinct role in DNA metabolism. The protein encoded by this gene is involved in excision repair and is located in both the mitochondria and nucleus, with translation initiation from the upstream start codon allowing for transport to the mitochondria and translation initiation from a downstream start codon allowing for transport to the nucleus. Additionally, alternate transcriptional splice variants, encoding different isoforms, have been characterized.

For Research Use Only