

# Recombinant Bovine Annexin V protein (His tag)

Catalog Number:PDEB100001



**Note:** Centrifuge before opening to ensure complete recovery of vial contents.


## Description

<b>Synonyms</b>	ANXA5;Annexin V;Unconjugated Annexin V;Annexin-5;Anchorin CII;Calphobindin I;Endonexin II;Lipocortin V;Placental anticoagulant protein 4;Thromboplastin inhibitor;Vascular anticoagulant-alpha
<b>Species</b>	Bovine
<b>Expression Host</b>	E.coli
<b>Sequence</b>	Met 1-Asp 320
<b>Accession</b>	P81287
<b>Calculated Molecular Weight</b>	35.1 kDa
<b>Observed molecular weight</b>	35 kDa
<b>Tag</b>	N-His

## Properties

<b>Purity</b>	> 95 % as determined by reducing SDS-PAGE.
<b>Endotoxin</b>	Please contact us for more information.
<b>Storage</b>	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
<b>Shipping</b>	This product is provided as lyophilized powder which is shipped with ice packs.
<b>Formulation</b>	Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
<b>Reconstitution</b>	Please refer to the printed manual for detailed information.

## Data



> 95 % as determined by reducing SDS-PAGE.

## Background

Peptidyl-prolyl cis-trans isomerase A, also known as PPIase A, Rotamase A, Cyclophilin A, Cyclosporin A-binding protein, PPIA and CYPA, is a cytoplasm protein which belongs to the cyclophilin-type PPIase family and PPIase A subfamily. Cyclophilins (Cyps) are a family of proteins found in organisms ranging from prokaryotes to humans. These molecules exhibit peptidyl-prolyl isomerase activity, suggesting that they influence the conformation of proteins in cells. PPIA / Cyclophilin A accelerate the folding of proteins. It catalyzes the cis-trans isomerization of proline imidic peptide bonds in oligopeptides. PPIA / Cyclophilin A is secreted by vascular smooth muscle cells in response to inflammatory

## For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623

Web: [www.elabscience.com](http://www.elabscience.com)

Tel: 1-832-243-6086

Email: [techsupport@elabscience.com](mailto:techsupport@elabscience.com)

Fax: 1-832-243-6017

# Recombinant Bovine Annexin V protein (His tag)

Catalog Number:PDEB100001



stimuli, and could thus contribute to atherosclerosis. It is not essential for mammalian cell viability. PPIA / Cyclophilin A can interact with several HIV proteins, including p55 gag, Vpr, and capsid protein, and has been shown to be necessary for the formation of infectious HIV virions.

## For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623

Web: [www.elabscience.com](http://www.elabscience.com)

Tel: 1-832-243-6086

Email: [techsupport@elabscience.com](mailto:techsupport@elabscience.com)

Fax: 1-832-243-6017