



A Reliable Research Partner in Life Science and Medicine

Purified Anti-Human CD273 Antibody[24F.10C12]

catalog number: E-AB-F11750P

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Reactivity Human

Immunogen Recombinant Human CD273 protein

Host Mouse

 Isotype
 Mouse IgG2a, κ

 Clone
 24F.10C12

Purification >98%, Protein A/G purified


Buffer Phosphate-buffered solution, pH 7.2, containing 0.05% non-protein stabilizer.

Dialyze to completely remove the stabilizer prior to labeling.

Applications Recommended Dilution

FCM 2 μ g/mL(0.5×10⁶-1×10⁶ cells)

Data



HEK293T cells transfected with pcDNA3.1 plasmid encoding Human CD273 gene were stained with 0.2 μg Purified Anti-Human CD273 Antibody[24F.10C12] (Right) and 0.2 μg Mouse IgG2a, κ Isotype Control (Left), followed by Elab

Fluor[®] 647-conjugated Goat Anti-Mouse IgG Secondary Antibody.

Preparation & Storage

Storage Storage Store at 4°C valid for 12 months or -20°C valid for long term storage, avoid freeze /

thaw cycles.

Shipping Ice bag

Background

T cells require a signal induced by the engagement of the T cell receptor and a "co-stimulatory" signal(s) through distinct T cell surface molecules for optimal T cell activation and tolerance. Members of the B7 superfamily of counter-receptors were identified by their ability to interact with co-stimulatory molecules found on the surface of T cells. Members of the B7 superfamily include B7-1 (CD80), B7-2 (CD86), B7-H1 (PD-L1), B7-H2 (B7RP-1), B7-H3, and PD-L2 (B7-DC).

For Research Use Only

 Toll-free: 1-888-852-8623
 Tel: 1-832-243-6086
 Fax: 1-832-243-6017

 Web: www.elabscience.com
 Email: techsupport@elabscience.com
 Rev. V1.6