
Recombinant mouse TGF-beta 2/TGFB2 protein (His Tag)

Catalog Number: PDEM100291

Note: Centrifuge before opening to ensure complete recovery of vial contents.

SpeciesMouseSourceE.coli-derived Mouse TGF-beta 2 protein Leu21-Arg302, with an N-terminaCalculated MW30.9 kDaObserved MW38 kDaAccessionP27090Bio-activityNot validated for activityPropertiesPurity> 95% as determined by reducing SDS-PAGE.Endotoxin< 10 EU/mg of the protein as determined by the LAL method	
Calculated MW30.9 kDaObserved MW38 kDaAccessionP27090Bio-activityNot validated for activityPropertiesPurity>95% as determined by reducing SDS-PAGE.	
Observed MW38 kDaAccessionP27090Bio-activityNot validated for activityProperties>95% as determined by reducing SDS-PAGE.	l His
AccessionP27090Bio-activityNot validated for activityPropertiesPurity> 95% as determined by reducing SDS-PAGE.	
Bio-activityNot validated for activityPropertiesPurity>95% as determined by reducing SDS-PAGE.	
Properties Purity > 95% as determined by reducing SDS-PAGE.	
Purity > 95% as determined by reducing SDS-PAGE.	
· ·	
Endotoxin < 10 EU/mg of the protein as determined by the LAL method	
Storage Generally, lyophilized proteins are stable for up to 12 months when stored	at -20 to -80
°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Alic	uots of
reconstituted samples are stable at $< -20^{\circ}$ C for 3 months.	
Shipping This product is provided as lyophilized powder which is shipped with ice p	oacks.
Formulation Lyophilized from a 0.2 µm filtered solution in PBS with 5% Trehalose and	5%
Mannitol.	
Reconstitution It is recommended that sterile water be added to the vial to prepare a stock	solution of
0.5 mg/mL. Concentration is measured by UV-Vis.	





> 95 % as determined by reducing SDS-PAGE.

Background

Transforming growth factor beta 2 (TGF- β 2) is a member of TGF-beta superfamily that shares a characteristic cysteine knot structure. Mice with TGF- β 2 gene deletion show defects in development of cardiac, lung, craniofacial, limb, spinal column, eye, inner ear and urogenital systems. All TGF- β isoforms signal via the same heteromeric receptor complex, consisting of a ligand binding TGF- β receptor type II (T β R-II), and a TGF- β receptor type I (T β R-I). Signal transduction from the receptor to the nucleus is mediated via SMADs. TGF- β expression is found in cartilage, bone, teeth, muscle, heart, blood vessels, haematopoitic cells, lung, kidney, gut, liver, eye, ear, skin, and the nervous system.