Elabscience®

RPS6KA5 Polyclonal Antibody

catalog number: E-AB-92667

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description	
Reactivity	Human;Mouse;Rat
Immunogen	A synthetic peptide of human RPS6KA5
Host	Rabbit
Is otype	IgG
Purification	Affinity purification
Buffer	Phosphate buffered solution, pH 7.4, containing 0.05% stabilizer and 50% glycerol.

Applications	Recommended Dilution
WB	1:500-1:2000

Data



Western blot analysis of C6 using RPS6KA5 Polyclonal

Antibody at 1:500 dilution.

Observed-MV:90 kDa

Calculated-MV:61 kDa/81 kDa/89 kDa

Preparation & Storage Storage Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles. Shipping The product is shipped with ice pack,upon receipt,store it immediately at the temperature recommended.

Background

Serine/threonine-protein kinase that is required for the mitogen or stress-induced phosphorylation of the transcription factors CREB1 and ATF1 and for the regulation of the transcription factors RELA, STAT3 and ETV1/ER81, and that contributes to gene activation by histone phosphorylation and functions in the regulation of inflammatory genes. Phosphorylates CREB1 and ATF1 in response to mitogenic or stress stimuli such as UV-C irradiation, epidermal growth factor (EGF and anisomycin. Plays an essential role in the control of RELA transcriptional activity in response to TNF and upon glucocorticoid, associates in the cytoplasm with the glucocorticoid receptor NR3C1 and contributes to RELA inhibition and repression of inflammatory gene expression. In skeletal myoblasts is required for phosphorylation of RELA at 'Ser-276' during oxidative stress. In erythropoietin-stimulated cells, is necessary for the 'Ser-727' phosphorylation of STAT3 and regulation of its transcriptional potential. Phosphorylates ETV1/ER81 at 'Ser-191' and ' Ser-216', and thereby regulates its ability to stimulate transcription, which may be important during development and breast tumor formation. Directly represses transcription via phosphorylation of 'Ser-1' of histone H2A. Phosphorylates ' Ser-10' of histone H3 in response to mitogenics, stress stimuli and ECF, which results in the transcriptional activation of several immediate early genes, including proto-oncogenes c-fos/FOS and c-jun/JUN.

For Research Use Only

Toll-free: 1-888-852-8623 Web:<u>w w .elabscience.com</u>