

Recombinant Rat IL-5 protein (His Tag)

Catalog Number: PDMR100026

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Species	Rat
Source	HEK293 Cells-derived Rat IL-5 protein Met1-Val132, with an C-terminal His
Calculated MW	14.41 kDa
Observed MW	20-28 kDa
Accession	Q08125
Bio-activity	Not validated for activity

Properties

Purity	> 90% as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU/mg of the protein as determined by the LAL method
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80 °C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from a 0.2 µm filtered solution in PBS with 5% Trehalose and 5% Mannitol.
Reconstitution	It is recommended that sterile water be added to the vial to prepare a stock solution of 0.5 mg/mL. Concentration is measured by UV-Vis.

Data



SDS-PAGE analysis of Rat IL-5 proteins, 2µg/lane of Recombinant Rat IL-5 proteins was resolved with SDS-PAGE under reducing conditions, showing bands at 20-28 KD.

Background

For Research Use Only

Interleukin 5 (IL-5) is a member of the interleukin family. Interleukins are a group of cytokines (secreted proteins/ signaling molecules) that were first seen to be expressed by white blood cells (leukocytes) and has been found in a wide variety of body cells. Interleukin 5 or IL-5 is produced by T helper-2 cells and mast cells. It helps to stimulate B cell growth and increase immunoglobulin secretion and is considered a key mediator in eosinophil activation. Interleukin 5 (IL-5) has long been associated with several allergic diseases, including allergic rhinitis and asthma. Growth in the number of circulating, airway tissue, and induced sputum eosinophils have been observed in patients with these diseases. IL-5 also had something with the terminally differentiated granulocyte eosinophils. IL-5 was originally found as an eosinophil colony-stimulating factor. It has been proved to be a major regulator of eosinophil accumulation in tissues and can modulate eosinophil behavior at every stage from maturation to survival.