

A Reliable Research Partner in Life Science and Medicine

APC Anti-Mouse CD40 Antibody[FGK4.5/FGK45]

Catalog Number: E-AB-F1028UE

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Reactivity Mouse Rat Host

Isotype Rat IgG2a, ĸ FGK4.5/FGK45 Clone No.

APC Rat IgG2a, κ Isotype Control[2A3] [Product E-AB-F09833E] Isotype Control

Conjugation

Conjugation Information APC is designed to be excited by the Red (627-640 nm) laser and detected using an

optical filter centered near 660 nm (e.g., a 660/20 nm bandpass filter).

Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer. Storage Buffer


Applications Recommended usage

FCM

Each lot of this antibody is quality control tested by flow cytometric analysis. Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use. We suggest each investigator should titrate the reagent to obtain optimal results [The recommended concentration is 0.1-1 μg/10⁶ cells

in 100 µL volume].

Data



C57BL/6 murine splenocytes are stained with APC Anti-Mouse CD40 Antibody (filled gray histogram). Unstained splenocytes (empty black histogram) are used as control.

Preparation & Storage

Keep as concentrated solution. Storage

This product can be stored at 2-8°C for 12 months. Please protected from prolonged

exposure to light and do not freeze.

Shipping Ice bag

Antigen Information

Alternate Names B-cell surface antigen CD40;Bp50;CD40;CD40L receptor;Cd40;Tnfrsf5;Tumor necrosis

factor receptor superfamily member 5

Uniprot ID P27512 Gene ID 21939

For Research Use Only

Tel: 1-832-243-6086 Fax: 1-832-243-6017 Toll-free: 1-888-852-8623 Web:www.elabscience.com

Rev. V1.5

Elabscience Bionovation Inc.



A Reliable Research Partner in Life Science and Medicine

Background

CD40 is a 48 kD type I transmembrane glycoprotein also known as Bp50. It is a member of the tumor necrosis factor receptor (TNFR) superfamily and is expressed on B cells, basal epithelial cells, macrophages, follicular dendritic cells, endothelial cells, and a subset of CD34+ hematopoietic progenitors. CD40 regulates B cell developmen t/maturation, Ig isotype switching and, in combination with other signals such as IL-4, protects B cells from surface lg-induced apoptosis and promotes proliferation. Interaction of CD40 with its ligand CD154 (gp39), which is expressed on activated T cells, is important in costimulation and immune regulation.

Fax: 1-832-243-6017 Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Web:www.elabscience.com